

# Isolation of *MA-ACS* Gene Family and Expression Study of *MA-ACSI* Gene in *Musa acuminata* Cultivar Pisang Ambon Lumut

LISTYA UTAMI KARMAWAN\*, SONY SUHANDONO, FENNY MARTHA DWIVANY

School of Life Sciences and Technology, Institut Teknologi Bandung, Jalan Ganesha No. 10, Bandung 40132, Indonesia

Received January 8, 2009/Accepted March 30, 2009

*Musa acuminata* cultivar pisang ambon lumut is a native climacteric fruit from Indonesia. Climacteric fruit ripening process is triggered by the gaseous plant hormone ethylene. The rate limiting enzyme involved in ethylene biosynthesis is ACC synthase (ACS) which is encoded by ACS gene family. The objective of this study is to identify *MA-ACS* gene family in *M. acuminata* cultivar pisang ambon lumut and to study the *MA-ACSI* gene expression. The result showed that there were nine *M. acuminata* ACS gene family members called *MA-ACSI*-9. Two of them (*MA-ACSI* and *MA-ACS2*) were assessed using reverse transcriptase PCR (RT-PCR) for gene expression study and it was only *MA-ACSI* correlated with fruit ripening. The *MA-ACSI* gene fragment has been successfully isolated and characterized and it has three introns, four exons, and one stop codon. It also shows highest homology with *MACSI* gene from *M. acuminata* cultivar Hsian Jien Chiao (GenBank accession number AF056164). Expression analysis of *MA-ACSI* using quantitative PCR (qPCR) showed that *MA-ACSI* gene expression increased significantly in the third day, reached maximum at the fifth day, and then decreased in the seventh day after harvesting. The qPCR expression analysis result correlated with the result of physical analysis during fruit ripening.

Key words: pisang ambon lumut, *MA-ACS* gene family, gene characterization, gene expression, quantitative PCR (qPCR)

## INTRODUCTION

*Musa acuminata* cultivar pisang ambon lumut is one of thousands existing banana cultivars around the world. It is well known in Indonesia and called pisang ambon lumut because it shows yellowish green peel when it has ripen (Verheij & Coronel 1992) which resembles the moss color ('lumut' is moss in Indonesia). Pisang ambon lumut has delicious sweet taste, specific aroma, and texture. Since banana (*Musa* sp.) is the world's fourth primary food after rice, wheat, and corn (Arias *et al.* 2003), pisang ambon lumut would become a promising export commodity in the future. Unfortunately, banana is classified into climacteric fruit which shows ripening process in a relative short period. This phenomenon becomes a disadvantage in commercial distribution of climacteric fruits, especially if it distributed in a long distance or if an unwanted delayed transport happens.

Manipulation strategy for delaying pisang ambon lumut would be advantageous for distance distribution; therefore a comprehensive data of pisang ambon lumut ripening process should be elucidated to define the manipulation technology. Previous study of pisang ambon lumut ripening process have been done by our group, such as protein isolation and characterization during banana ripening process (Insani 2006), isolation and characterization of ACS and ACO gene which involved in ethylene biosynthesis (Dwivany *et al.* unpublished data), and molecular strategy for controlling banana ripening using RNAi mechanism which silence specific genes involved in ethylene biosynthetic pathway (Dwivany *et al.* unpublished data).

Fruit ripening process is regulated by the plant gaseous hormone, ethylene. Biochemically, fruit ripening can be defined as the summation of changes in tissue metabolism that renders the fruit attractive for consumption by organisms that assist in seed release and dispersal. In spite of fruit ripening, ethylene has the ability to switch on and off hundreds genes affecting various process. The expression or suppression of these genes induce morphological, physiological, and biochemical changes in plants in diverse events such as seed germination, abscission, fruit ripening, senescence, and various kinds of biotic and abiotic stresses (Nath *et al.* 2006).

Ethylene is synthesized within the cell via the ethylene biosynthetic pathway. ACC synthase (ACS) which encoded by ACS gene is the rate limiting enzyme in ethylene biosynthesis (Tsuchisaka & Theologis 2004). It catalyzed the conversion of S-adenosyl-L-methionine (SAM) into 1-aminocyclopropane-1-carboxylic acid (ACC). In the next step, ACC is converted into ethylene, CO<sub>2</sub> and HCN by ACC oxidase (ACO) (Chaves & de Mello-Farias 2006).

In this study the gene family of ACC in pisang ambon lumut is isolated. The specific *M. acuminata* ACS (*MA-ACS*) gene family member which is involved in pisang ambon lumut ripening process has been further characterized. Expression analysis using quantitative PCR (qPCR) has been performed to quantify the expression of *MA-ACSI* gene during ripening process.

## MATERIALS AND METHODS

**Sampling.** Mature *M. acuminata* cultivar pisang ambon lumut were harvested after 12 weeks since the male flower emerged. Fingers of the second bunch (hand) of this banana were used for experiment to reduce data variation (Dadzie &

\*Corresponding author. Phone: +62-222-511157, Fax: +62-222-514307, E-mail: fenny@sith.itb.ac.id

Orchard 1997). This second bunch was stored in the room temperature and ripened naturally. Physical assessment and RNA isolation were conducted every two days in seven days since fruits were harvested (duplo). On the other hand, DNA isolate from banana plantlet leaf was obtained from Sadjuri (School of Life Sciences and Technology, Institut Teknologi Bandung).

**RNA Isolation and First Strand cDNA Synthesis from Banana Pulp.** RNA was isolated using Asif's method (Asif *et al.* 2000) from banana pulp. RNA is then converted into first strand cDNA using *SuperScript™ First-Strand Synthesis System for RT-PCR* from Invitrogen®. RNA was stored in -80 °C before used for cDNA fragment isolation and expression analysis.

**MA-ACS Gene Family Identification.** Nine pairs of specific primers for banana *MA-ACS* gene family (called ACS1rt F/R and ACS2 F/R – ACS9 F/R, Table 1) were designed based on *ACS* gene family in *M. acuminata* cultivar Hsian Jien Chiao (Huang *et al.* 2006). These primers are used for PCR analysis to isolate *ACS* gene family fragments in *M. acuminata* cultivar pisang ambon lumut. The resulting fragments which ranged between 102-346 bp were cloned into pGEM®-T Easy cloning vector, sequenced and compared with GenBank database using BLAST software ([www.ncbi.nlm.nih.gov/BLAST](http://www.ncbi.nlm.nih.gov/BLAST)).

**Expression Analysis of MA-ACS1 and MA-ACS2.** Total cDNA from unripen and ripen banana were used for qualitative expression analysis for *MA-ACS1* and *MA-ACS2* gene by PCR detection method using ACS1rt F/R and ACS2 F/R primer pairs. Furthermore, qPCR (quantitative Polymerase Chain Reaction) was performed for *MA-ACS1* using iQ™5 *Real-Time PCR Detection Systems* with iQ™ SYBR® Green Supermix reagent. The qPCR analysis was performed for *MA-ACS1* gene expression in the fruit pulp on first, third, fifth, and seventh day after harvesting. The ACS1rt F/R were specifically designed for qPCR and then assessed by melting curve analysis. Good qPCR primers will give single peak when were assessed with melting curve analysis (Dorak 2006).

Table 1. Oligonucleotides used in gene cloning reactions and qPCR assay

Primer	Sequence (5'-3')
ACS1F	TACGGCGAGGAGCACCCAAAT
ACS1R	CGATGTTTCAGGTGGCGGCTT
ACS1rtF	CCGAGACTGGATGAAGAAGAA
ACS1rtR	GTCTGGGTCAAATCTGGCTC
ACS2F	CTTGAGAACCATCCCGACC
ACS2R	TCATTGGCAGAAGTAGCACC
ACS3F	GGTACACCGTGGACGACAC
ACS3R	AGCCACGACTCGATGAGAT
ACS4F	GCTTCTCCTCATTCGTCATTC
ACS4R	GAGGTCAGGGTGGTCTTCAA
ACS5F	GCCAAATATGCCTTCCGTA
ACS5R	GTCTTAAACTCGGAAACGC
ACS6F	TTTAGCCTTCTCAGGTTTCA
ACS6R	AGCATTCTCTGAAGCTCGA
ACS7F	TTTCTCTGCTGACTGTGC
ACS7R	AGCCGGAGATTCCACATC
ACS8F	ATTACTTTCTCTGCTGCTGC
ACS8R	ACTCTGAAACGCCCTCCTT
ACS9F	GTGGATCGCCATCATCTTT
ACS9R	AGCCACGACTCGATGAGAT

Quantification of serial *MA-ACS1* gene expression was carried out using absolute quantification method (Livak & Schmittgen 2006). Calibration curve for the absolute quantification was made using serial dilution of plasmid containing *MA-ACS1* gene with known copy number for 10 times dilution, started from 300,000 until 30 plasmid copies. The resulting qPCR data for each dilution is then plotted to cycle threshold number to construct a calibration curve. The serial quantification and calibration curve reaction were done simultaneously in qPCR machine to reduce variation.

**MA-ACS1 Gene Fragment Isolation and Characterization.** *MA-ACS1* genomic and cDNA fragment were amplified using *MA-ACS1* F/R primers (Wicaksono 2006) by PCR method from DNA and cDNA template. The forward primer was located nearby the start codon and the reverse primer the stop codon. The resulting fragments were then cloned into pGEM®-T Easy cloning vector and sequenced. The genomic and cDNA fragment sequences were then compared and characterized using bioinformatics software, such as BioEdit, BLAST (<http://ncbi.nlm.nih.gov/blast>), and GENSCAN 1.0 (<http://genes.mit.edu/GENSCAN.html>).

**Physical Assessment Method for Banana Ripening Process.** Peel color, peel thickness, pulp to peel weight ratio, qualitative starch content and qualitative sugar content were analyzed for physical assessment of banana ripening process according to banana routine post-harvest screening method (Dadzie & Orchard 1997). Peel color and peel thickness was documented with digital camera. Pulp weight and peel weight were measured by digital weight balance and then plotted into a graph. Qualitative starch content within banana pulp was performed using iodine test while qualitative sugar content of pulp juice was performed using Benedict test.

## RESULTS

**MA-ACS Gene Family Identification.** PCR of pisang ambon lumut DNA template using ACS1rt F/R and ACS2 F/R – ACS9 F/R produced nine fragments with size varying from 102 to 346 bp (Table 2). Each amplicon showed high homology with the correlated *ACS* gene family in *M. acuminata* cultivar Hsian Jien Chiao. BLASTN identity percentage varied between 97-99% with E-value varied from 1e-41 to 7e-174 (data not shown).

**Expression Study of MA-ACS1 and MA-ACS2.** Expression analysis of *MA-ACS1* and *MA-ACS2* using RT-PCR showed that only *MA-ACS1* was expressed during ripening. In other hand, *MA-ACS2* expression was not detected by

Tabel 2. *MA-ACS* gene family fragments of *M. acuminata* cultivar pisang ambon lumut

PCR primer	Gene	GenBank ACC number
ACS1rt F/R	MA-ACS1	GQ406065
ACS2 F/R	MA-ACS2	GQ406066
ACS3 F/R	MA-ACS3	GQ406067
ACS4 F/R	MA-ACS4	GQ406068
ACS5 F/R	MA-ACS5	GQ406069
ACS6 F/R	MA-ACS6	GQ406070
ACS7 F/R	MA-ACS7	GQ406071
ACS8 F/R	MA-ACS8	GQ406072
ACS9 F/R	MA-ACS9	GQ406073

electrophoresis analysis (Figure 1). Further study using qPCR absolute quantification for *MA-ACSI* showed that its gene expression increased significantly in the third day, reached maximum at the fifth day, and then decreased in the seventh day after harvesting (Figure 2). This absolute quantification was generated based on the calibration curve of *MA-ACSI* which has been created before (data not shown).

**Isolation and Characterization of *MA-ACSI*.** Genomic fragment of *MA-ACSI* (1818 bp, ACC number GQ396304) which were amplified using PCR primers ACS1F and R showed highest homology (99% similarities) with *MACSI* gene from *M. acuminata* cultivar Hsian Jien Chiao (GenBank ACC number AF056164). The *MA-ACSI* cDNA fragment (1459 bp, ACC number GQ406064) which was amplified using PCR primers ACS1F and R (Wicaksono 2006) was also showed highest homology with the same accession number. When the genomic and cDNA fragments were aligned, it was revealed that *MA-ACSI* from pisang ambon lumut has three introns, four exons, and one stop codon (Figure 3). The fragment did not have start codon because the forward primer's position was in the downstream of the start codon. The *MA-ACSI* gene structure was also confirmed using GENESCAN1.0 analysis and showed the same result as the alignment analysis (Figure 4).

**Physical Assessment Method for Banana Ripening Process.** Peel color changed during pisang ambon lumut ripening started from green at the first day and gradually changed into yellowish green with black spots at the seventh




Figure 1. Electroferogram of PCR and RT-PCR analysis of *MA-ACSI* and *MA-ACS2*. 1: *MA-ACSI* genomic (346 bp); 2: *MA-ACSI* cDNA (172 bp) from ripen pisang ambon lumut fruit; 3: *MA-ACS2* genomic (321 bp); 4: *MA-ACS2* cDNA (not detected) from ripen pisang ambon lumut fruit; 5: 100 bp DNA ladder.




Figure 2. Absolute quantification of *MA-ACSI* copy number using qPCR.

day (Figure 5a). Fruit peel became thinner and pulp to peel weight ratio increased (Figure 5b, 6), starch become degraded (Figure 5c) and sugar were accumulated (Figure 5d).

## DISCUSSION

PCR result from DNA template using specific primers for the *ACS* gene family showed that there were at least nine *ACS* genes identified in pisang ambon lumut. Sequencing and BLASTN analysis of each amplicon showed that the nine *MA-ACS* genes in *M. acuminata* cultivar pisang ambon lumut have a relative low E-value for homology. Identification and characterization of these *MA-ACS* gene family in *M. acuminata* cultivar pisang ambon lumut gives an opportunity to explore the promoter of each gene family member. Spatial and temporal expression study of these promoters would be important to make it useful for other application, for example expression of antigenic protein for producing edible vaccine. Further study must be done for gene family promoter utilizations.

According to the qualitative expression analysis of *MA-ACSI* and *MA-ACS2* by RT-PCR showed that only *MA-ACSI* was expressed during pisang ambon lumut ripening process. Subsequently, the *MA-ACSI* gene became an interesting subject for further analysis. The qPCR data showed that the expression of *MA-ACSI* increased and then decreased at the end of ripening. This data correlated with the physical assessment during its.

Physical assessment shows that chlorophyll was gradually degraded. Chlorophyll degradation is caused by the chlorophyllase activity which revealed the peel color from green into yellowish green (Chaves & de Mello-Farias 2006). Moreover, it was shown that starch within the pulp was also gradually degraded. Iodine test showed a decreasing of starch content, detected by decreased black color after stained with iodine reagent (Page 1981). Whilst, during ripening process the sugar content was gradually increased, detected by green color of Benedict reagent with yellow or red brick sediment (Page 1981). Starch degradation into sugar is catalyzed by  $\alpha$ -amylase activity. As the sugar is accumulated in the fruit pulp, it is become more osmotic than the peel (Dadzie & Orchard 1997). As shown in the result, the consequence of sugar accumulation in fruit pulp was the gradual decrease in fruit peel thickness because of water evaporation and water content movements from peel to pulp.

Chlorophyllase and  $\alpha$ -amylase activity which increased during the pisang ambon lumut ripening is influenced by ethylene (Chaves & de Mello-Farias 2006). Therefore, it was suggested that the increased expression of *MA-ACSI* correlates with ethylene production then induced the biochemical changes during fruit ripening. Decreased expression of *MA-ACSI* in the late ripening maybe a result from negative feedback regulation when ethylene accumulate in the late climacteric phase as mentioned by Inaba (2007).

Structural analysis of *MA-ACSI* gene was also performed. A longer fragment of *MA-ACSI* was isolated and characterized. The result showed that the *MA-ACSI* gene structure was similar to another *MA-ACS* gene family in other banana cultivar

```

TACGGCGAGGAGCCACCAATCAGCAGATCCCTCTCTCGGATCGCGACCACGACGGCCATGGCGGAACTCCTCTACTTCGATGGGTGGAAAGGCTACG 100
Y G E E H P N Q Q I L S R I A T N D G H G E N S S Y F D G W K A Y
AGAAGGATCCTTTCCACCTCACCGACAAACCCACGGGGTCCATCCAAATGGGACTCGCAGAAAACAGGgttagagttccctccatgggtgatgattaatcgc 200
E K D P F H L T D N P T G V I Q M G L A E N Q
acatgcctcccgccaattgccactccctggcgttgctaaatcctgtatgtgggttttgggctcttctctccctcagCTTTCCTCGGACTTGGTCCGAG 300
L S L D L I R
ACTGGATGAAGAAGAACCACAGGCTTCGATCTGCAACGAAAGGGGCTCAGAGTTCAAAGCAATTGCCAACTTTCAGGACTATCATGGCTCCGAGC 400
D W M K K N P Q A S I C T E E G V S E F K A I A N F Q D Y H G L P T
CTTCGAAAAGGtaaatgatctcaaccacaaacgcagcgtgagcgtgcttgctccactgtccaagtagctacatacgtccaatatgataaagctgggact 500
F R K
gacagccacttacggcccgagccctgcttgcctaccctggataaggataaagcctaatgatgggtgtgatcttgcgcaacgcgcaggCCATCGCCGAGTTCA 600
A I A Q F
TGGAGAAGGTGAGAGGGGACGAGCCAGATTTGACCCAGACCGCATCGTGTGAGCGGTGGAGCCACCGGGCTCAGGAAACCATCGCCTTTTGCCTGGC 700
M E K V R G G R A R F D P D R I V M S G G A T G A Q E T I A F C L A
TGATCCTGGCGAGGCTTCTTGATTCCAACGCCATATTATCGGGgtaagtaatttaggtgtactaaatctaccgagttctttatccggcagaggatcta 800
D P G E A F L I P T P Y Y P G
ggcatctgcatggtttccagATTGATCGAGACTTCAGGTGGAAGCAGGAGTTCAGCTCCTCCCATTCACCTGCCACAGTTCACCAAGATTCAAGATCA 900
F D R D F R W R T G V Q L L P I H C H S S N K F K I
CCCAAGCCGCACTGGAGACTGCTTACAGGAAGGCTCGAACTCACACATTAGAGTCAAAGGAATACTGGTGACCAACCCATCGAACCTCTGGGACACAC 1000
T Q A A L E T A Y R K A R N S H I R V K G I L V T N P S N P L G T T
CATGGACAGAGAGAGCTGAGAACCTAGTCACTTCGTCACGAGAAAAGGATGCACCTTGGTGTGGCAGGAGATCTTctcCGGAACCGTCTTCGACAG 1100
M D R E T L R T L V S F V N E K R M H L V C D E I F S G T V F D K
COGAGTTAAGTGAAGGTCCTCGAGGTTGATCGAAGACGAGCCCTACTGGCAGACGGGATCTGATTTCACATCGGCTACAGCCTCTCCAAGGACCTGGGCGTCC 1200
P S Y V S V S E V I E D E P Y C D R D L I H I A Y S L S K D L G V
CTGGCTTCGGCTGGGGTGCATATACTCCTACACGACGCCGTGGTCAAGTGCAGGAGGAAGATGTCGAGCTTTGGACTGGTCTCGTCCGACAGCAGCA 1300
P G F R V G V I Y S Y N D A V V S C A R K M S S F G L V S S Q T Q H
CCTGCTCGCTTCCATGTTGGGAGACGAGGAGTTCACCAAGGTTTCTTAGCGACGAGCCGAGCAGGTTGTGCGGGCGGCCAGGGTCTTTACGGACGGC 1400
L L A S M L G D E E F T T S F L A T S R T R L C G R R R V F T D G
CTCAAGCGAGTGGGATTGCTTGGACGGCAACCGGGGCTGTTCTGCTGGATGGACTTGAGGCCGTTGCTGAAGGAAGCGAGCGTGGAGCGGGAGC 1500
L K R V G I H C L D G N A G L F C W M D L R P L L K E A T V E A E
TCCGGCTGTGGCGGTTGATCATCAACGAGCSTGAAGCTCAACATCTCGCGGGGCTGTCCTTCCACTGCTCGGAGCGGGGTTGAGGTTGCTTCCG 1600
L R L W R V I I N D V K L N I S P G S S F H C S E P G W F R V C F A
CAACATGGACGACACGGCCATGAAGATAGCGCTGAGGAGGATCGAGAGTTTCTGCTGACCGGAGAACGACGCCCTGTGCAGGGGGAAGAACAAGAGGAG 1700
N M D D T A M K I A L R R I E S F V Y R E N D A A V Q A K N K R R
TGGGACGAAGCGCTCGGGCTGAGCTTGCCTCGTGGAGGTTGAGGATCCGACCATCATGACACCCACATCTGATGCTCCCACTCGCCTCTCGTTCAG 1800
W D E A L R L S L P R R R F E D P T I M T P H L M S P H S P L V Q
COGCCACCTGAaacatcg 1818
A A T *
    
```

Figure 3. Structure of the putative ACC synthase gene, MA-ACS1 (ACC number GQ396304), from *M. acuminata* cultivar pisang ambon lumut. Transcribed coding sequences are shown in uppercase letters, while intron, and UTR sequences are represented in lowercase letters. Stop codon are underlined.




Figure 4. Structure prediction of putative MA-ACS1 gene fragment using GENSCAN1.0 MA-ACS1 had four exons, three intron, and one stop codon.

and ACS gene family in another plant taxa such as *Arabidopsis* (van der Straeten *et al.* 1992) and tomato (Olson *et al.* 1991). It showed that the MA-ACS1 is conserved among plant species and may have a similar function.

There are still a lot of unexplored properties of pisang ambon lumut ripening process and comprehensive studies should be carried out to reveal them. In the future, biotechnology would help us to discover, to use, and to improve the biological potential of pisang ambon lumut for human welfare.

#### ACKNOWLEDGEMENT

This study is sponsored by PT. Indofood Sukses Makmur – Indofood Riset Nugraha 2006. We thank PT. Indofood Sukses Makmur for the research grant from “Indofood Riset Nugraha 2006”. We also thank to Sadjuri for pisang ambon lumut DNA sample.




Figure 5. Qualitative physical assessment during pisang ambon lumut ripening. Two samples were assessed every two days from day 1 until day 7 since harvested. a. Peel colour changes; b. transversal section of banana fruit, showed gradual decrease in peel thickness; c. Iodin test for starch detection, black stain showed detected starch; d. Benedict test for sugar detection.




Figure 6. Average ratio of banana pulp to peel weight during pisang ambon lumut ripening.

## REFERENCES

- Arias P, Dankers C, Liu P, Pilkauskas P. 2003. *The World Banana Economy, 1985-2002: FAO Commodity Studies*. Rome: FAO.
- Asif MA, Dhawan P, Nath P. 2000. A Simple Procedure for the Isolation of High Quality RNA from Ripening Banana Fruit. *Plant Mol Biol Rep* 18:109-115.
- Chaves ALS, de Mello-Farias PC. 2006. Ethylene and fruit ripening: from illumination gas to the control of gene expression, more than a century of discoveries. *Genet Mol Biol* 29:508-515.
- Dadzie BK, Orchard JE. 1997. *Routine Post-Harvest Screening of Banana/Plantain Hybrids: Criteria and Methods*. France: INIBAP.
- Dorak MT. 2006. *Real-time PCR*. New York: Taylor & Francis Group.
- Huang FC, Do YY, Huang PL. 2006. Genomic organization of a diverse ACC synthase gene family in banana and expression characteristics of the gene member involved in ripening of banana fruits. *J Agric Food Chem* 54:3859-3868.
- Inaba A. 2007. Studies on the internal feedback regulation of ethylene biosynthesis and signal transduction during fruit ripening, and the improvement of fruit quality. *J Japan Soc Hort Sci* 76:1-12.
- Insani M. 2006. *Isolasi dan Analisis SDS-Page Total Protein dari Pematangan Buah Pisang Ambon Lumut (Musa sp.)* [Thesis]. Bandung: Institut Teknologi Bandung.
- Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta CT}$  method. *Methods* 25:402-408.
- Nath P, Trivedi PK, Sane VA, Sane AP. 2006. Role of ethylene in fruit ripening. In: Khan NA (ed). *Ethylene Action in Plants*. Heidelberg: Springer-Verlag Berlin Heidelberg.
- Olson CD, White JA, Edelman L, Harkins RN, Kende H. 1991. Differential expression of two genes for 1-aminocyclopropane-1-carboxylate synthase in tomato fruits. *Proc Natl Acad Sci* 88:5340-5344.
- Page DS. 1981. *Principles of Biological Chemistry*. 2<sup>nd</sup> ed. Jakarta: Erlangga.
- Tsuchisaka A, Theologis A. 2004. Unique and overlapping expression patterns among the arabidopsis 1-Amino-cyclopropane-1-carboxylate synthase gene family members. *Plant Physiol* 136:2982-3000.
- van der Straeten D *et al.* 1992. Cloning genetic mapping, and expression analysis of an *Arabidopsis thaliana* gene that encodes 1-aminocyclopropane-1-carboxylate synthase. *Proc Natl Acad Sci* 89:9969-9973.
- Verheij EWM, Coronel RE. 1992. *Plant Resource of South East Asia: Edible Fruits and Nuts*. Volume 2. Bogor: PROSEA Foundation.
- Wicaksono BD. 2006. *Isolasi, Kloning, dan Analisis Ekspresi Gen ACC Sintase (ACS) dari Buah Pisang Ambon* [Thesis]. Bandung: Institut Teknologi Bandung.