

# Prediction of Physicochemical Properties of Indonesian Indica Rice Using Molecular Markers

PUJI LESTARI<sup>1,2</sup>, REFLINUR<sup>1,2</sup>, HEE-JONG KOH<sup>2\*</sup>

<sup>1</sup>*Indonesian Center for Agricultural Biotechnology and Genetic Resources Research and Development, IAARD  
Jalan Tentara Pelajar No. 3A, Bogor 16111, Indonesia*

<sup>2</sup>*College of Agriculture and Life Science, Seoul National University, San56-1, Kwanak-gu, Seoul, South Korea*

Received October 28, 2013/Accepted May 8, 2014

Physicochemical properties determine the palatability and cooking quality of rice, which must be determined efficiently in order to satisfy consumer demand. To date, little information exists on the use of molecular markers to predict physicochemical properties of the “indica” rice varieties found in Indonesia. The objective of this study was to investigate physicochemical properties and genetic variation of Indonesian rice varieties, and to formulate regression equations to analyze sets of DNA markers which could predict amylose content (AC), protein content (PC) and pasting properties of the varieties. A total of 24 Indonesian indica rice varieties were chosen based on their genetic background and agricultural characteristics. We then measured selected physicochemical properties, and genotyped the varieties using 30 DNA markers. The chosen varieties showed favorable values for PC, AC, and six rapid viscosity analyzer (RVA) pasting properties, which was supported by molecular data. As demonstrated by principal component analysis (PCA), markers could provide a complementary method for differentiating rice varieties, as an alternative to measuring physicochemical properties. PCA analysis also allowed us to establish marker sets using multiple regression analysis. We formulated eight model regression equations comprising data regarding 15 to 19 markers with high coefficients ( $R^2=0.98-0.99$ ). The formulas provided results that consistently correlated and therefore predicted the physicochemical properties of indica rice. Further validation of these marker sets may provide rapid and efficient means for predicting the physicochemical properties of Indonesian-bred indica rice in the future.

Keywords: indica rice, molecular marker, physicochemical properties, multiple regression

## INTRODUCTION

Rice is one of the major cereal crops in the world and has been bred for many years. As a consequence, it has undergone many genetic changes before achieving its current form. In Asia, indica and japonica types are most commonly used, but they differ in physical and chemical properties. These differences have led to differentiation of food habits in different regions, japonica rice denotes short grain cultivars, with a soft and sticky texture; while indica rice is typically long-grain and remains separate, light and fluffy after cooked. There are many types of indica rice, which are usually grown in hot climates, particularly in Southern and Southeast Asian countries such as India, Thailand, Vietnam and Indonesia (Izawa 2008; Mohapatra & Bal 2011). Tropical climate conditions in Indonesia along with consumer taste, has led to a preference for growing indica rice over other types.

Eating and/or cooking quality is determined primarily by the physicochemical properties of rice starch, and such characteristics have been given much research attention (Ramesh *et al.* 1999; Kuo *et*

*al.* 2001). In evaluating rice grain quality, amylose content (AC) and protein content (PC) are among the important factors determining eating quality (palatability) such as texture, flavor and stickiness; and cooking qualities (such as grain elongation, gel consistency, and gelatinization temperature). However, rice varieties with similar AC may have different starch pasting viscosity. Higher AC and different levels of PC and pasting properties of indica varieties are observed in comparison with japonica rice. Genetic studies have been increasingly conducted in relation to physicochemical properties with respect to indica rice palatability/cooking quality. The most current discoveries elucidated from genetic level research has been integrated with conventional breeding in order to develop of new rice varieties (Pooni *et al.* 1993; Li *et al.* 2003; Bao *et al.* 2008).

Conventional methods have commonly been used to measure the physicochemical properties of rice to inform efforts to enhance the palatability and cooking quality of rice varieties. PC and AC are typically measured based on Association of Official Analytical Chemists (AOAC) calculations using specific equipment. Pasting viscosity can be estimated using a rapid viscosity analyzer (RVA)

\*Corresponding author. Phone: +82-2-8804541/4978,  
Fax: +82-2-8732056, E-mail: heejkoh@snu.ac.kr

and amylograph. All these measurements require large rice samples, and technically it is a difficult, inefficient, costly and time-consuming endeavor. To provide indica rice cultivars which can satisfy consumers with respect to consumption quality, it is necessary to establish a rapid detection method to enable efficient selection.

Advances in scientific understanding of molecular markers confirmed their potential application to genetic research of different kinds, and for molecular-assisted breeding (Tian *et al.* 2005). Several types of DNA markers such as simple sequence repeat (SSR), single nucleotide polymorphism (SNP), sequence tagged-site (STS), and insertions/deletions (Indels) related to the physicochemical properties of rice have been identified. These markers can be used for genotyping, genetic diversity analyses and genetic mapping. DNA markers have also been used to assist in breeding programs aimed at improving the eating quality of rice, including predicting specific food qualities and physicochemical properties of japonica rice (Ohtsubo *et al.* 2002, 2003; Lestari *et al.* 2009). However, no study has been published regarding detection of physicochemical properties of Indonesian indica rice using DNA markers. Most markers for physicochemical properties determining palatability/cooking quality have been developed based on polymorphic and association analyses (Bao *et al.* 2006a,b; Patindol *et al.* 2010; Zhao *et al.* 2012). By using molecular markers developed for japonica and indica genomes, predicting physicochemical properties of indica rice should be much more efficient.

In this study, 24 indica rice varieties originating from Indonesia were analyzed for their physicochemical properties, and genotyped using 30 DNA markers. Genetic variation among the rice varieties was investigated and then model equations were formulated using the molecular markers in order to predict the physicochemical properties of Indonesian indica rice.

## MATERIALS AND METHODS

**Plant Materials and DNA Extraction.** For this study, two landraces and 22 indica rice varieties/strains bred in Indonesia were chosen from the germplasm collection of Indonesian Center for Agricultural Biotechnology and Genetic Resources Research and Development (Table 1). Information on their genetic background and known agricultural characteristics was also collected. The plant samples used for physicochemical analysis, came from all varieties and were grown using conventional cultural

practices at the experimental farm of Seoul National University, Korea. Rice grains were dried to 15% moisture content before analysis. For DNA isolation and analysis, plants for each rice variety were grown in a greenhouse until tillering stage. Genomic DNA was extracted from young leaf tissue using the cetyl trimethyl ammonium bromide (CTAB) method as described by Murray and Thompson (1980).

**Primer and PCR Amplification.** For DNA analysis in this study, we used a total of 30 molecular markers comprising SSR, STS, dCAPS/CAPS, and Indel. Some of these had been reported previously (Lestari *et al.* 2009) and some were newly developed during the study. The list of markers and their sequences is presented in Table 2. Standard PCR was carried out in a PTC-200 Peltier Thermal Cycler (MJ Research, Inc.) under the following conditions: a total of 35 cycles comprising 1 minute at 95 °C, then 30 seconds at 55 °C and finally, 1 min at 72 °C. PCR amplification of DNA was performed in a total volume of 20 µL with the following PCR reagents: 2 µL of DNA at 20 ng/µL, 2 µL of 10X buffer containing 25 mM MgCl<sub>2</sub>, 1 µL of 2.5 mM dNTPs, 1 U of Taq Polymerase (Intron Biotechnology, Korea), and 1 µL each of forward and reverse primers (10 µM). The amplicons were separated by electrophoresis on 3% agarose gels and/or by non-denaturing 8% polyacrylamide gels stained with ethidium bromide (Model MG, CBG Scientific Co.).

**Measurement of Physicochemical Properties.** The rice grains were hulled and milled to 91% yield. Protein content (PC) was calculated by means of multiplication of total nitrogen by 5.95 after determination of the nitrogen content of rice material using the Micro-Kjeldahl method (AOAC 1995). Amylose content (AC) of milled rice was measured by using the relative absorbancy of starch-iodine color in solution of 100-mesh rice flour digested according to the previous method (Perez & Juliano 1978). Pasting properties were determined on a rapid visco analyser (RVA) following the procedures listed in the manufacturer user manual (NewPort Sci. Co., Australia). The pasting profiles of rice starch samples were described based on six parameters: peak viscosity (PV), hot paste viscosity (HPV), cool paste viscosity (CPV), breakdown viscosity (BDV=PV-HPV), setback viscosity (SBV=CPV-PV), and consistency viscosity (CTV=CPV-HPV) according to Bao and Xia (1999). These rice starch properties were measured and confirmed in three replications.

**Data Analysis.** PCR amplicon of each marker was converted to binary values denoting the base

Table 1. Twenty four Indonesian-bred rice varieties used in this study and their characteristics

Name	Variety	Pedigree	Type	Days to maturity	Yield potency (ton/ha)	Cooked rice texture	Amylose content (%)*	Unhulled rice grain color
Rojolele	landrace		Hairy grain surface, lowland	155	4.2	aromatic, fluffy	medium	yellow
Kalimutu	landrace		cereh	90-95	2.5	medium	medium	clear yellow
Jatiluhur	variety	Tox1011/Ranau	cereh, upland	110-115	3.5	separate	high	dark yellow
Fatmawati	variety	BP68C-MR-4-3-2/Maros	cereh, new type	105-115	9	good, fluffy	medium	clear yellow
Batang Piaman	variety	IR25393-57/RD203//IR27316-96//SPLR7735/ SPLR2792	cereh, lowland	100-117	7.6	separate	high	clear yellow
Ciliwung	variety	IR38//2*Pelita 1-1//IR4744-128-4-1-2	cereh, lowland	117-125	6.5	fluffy	medium	clear yellow
Angke	variety	IR64*6//IRBB5	cereh, lowland	110-120	7.5	fluffy	medium	clear yellow
Konawe	variety	S487B-75/2*IR19661-131-3-1//2*IR64	cereh, lowland	110-120	8	fluffy	medium	clear yellow
Memberamo	variety	B6555B-199-40/Barumun	cereh, lowland	115-120	7.5	fluffy	low	clear yellow
Cirata	variety	IR9129-159-3//IR5975	cereh, upland	115-125	6.5	fluffy	medium	yellow, purple on tip
Batanghari	variety	Cisadane//IR19661-131-1-3-1-3	cereh, tidal swamp	122-128	6.5	separate	high	clear yellow
Ciherang	variety	IR18349-53-1-3-1-3//3*IR19661-131-3-1-3//4*IR64	cereh, lowland	116-125	8.5	fluffy	medium	clear yellow
Cimelati	variety	Memberamo//IR66160/Memberamo	cereh, semi new type	118-125	7.5	good, fluffy	low	clear yellow
Cigeulis	variety	Ciliwung/Cikapundung//IR64	cereh, lowland	115-125	8	fluffy	medium	clear yellow
Conde	variety	IR64*6//IRBB7	cereh, lowland	115-125	7.5	fluffy	medium	clear yellow
Singkil	variety	IR35432-33-2//IR19661-131-3-1//Ciliwung//IR64	cereh, lowland	115-125	7	good, fluffy	medium	clear yellow
Batang Gadis	variety	IR64/NDR308//IR64	cereh, lowland	108-112	7.5	fluffy	medium	clear yellow
Pepe	variety	Simariti/4*IR64	cereh, lowland	120-128	8.1	fluffy	medium	yellow
Maros	variety	Markoti//IR64	cereh, lowland	110-128	9	fluffy	medium	clear yellow
Tukad Balian	variety	IR48613-54-3-3-1//IR28239-94-2-3-6-2	cereh, sometime hairy, lowland	105-115	7	fluffy	medium	straw-like yellow
Cisokan	variety	PB36/Pelita 1-1	cereh, sometime hairy, lowland	110-120	6	separate	high	clear yellow
Cibodas	variety	B7004D-MR-10-1//B6992F-MR-26	cereh, lowland	117-126	7	medium	medium	clear yellow
Logawa	variety	Cisadane/Bogowonto//2*Cisadane	cereh, lowland	110-120	7.5	separate	high	yellow
Sintanur	variety	Lusi//B7136C-MR-22-1-5 (Bengawan Solo)	cereh, lowland	115-125	7	fluffy	low	clear yellow

\*the type of AC was grouped as low (&lt;20%), medium (20-25%), and high (&gt; 25%) according to Allidawati and Bambang (1989).

Table 2. List of 30 primers and their sequences used in this study

Primer	Sequences	
	Forward (‘5—3’)	Reverse (‘5—3’)
A7	TGCCTCGCACCAGAAATAG	TGCCTCGCACCATGAG
E30	TACCTGGTTGATGTATACAGATCTGGTT	ATCCCTCGATCCCTCTAGCATTAT
F4	ACCACTCCATATATATCATCCAAAG	ACCACTCCATATACCACAAGG
G4	GAGACCGATATGCGATTC	GTGGTGTTTAGATCCAGAGACTTA
G28	GGCGGTTCGTTCTGCGAT	GGAGAATCCCACAGTAAGTTTTCTTTG
S13	GTCGTTCCCTGTGGTTAGGACAGGGT	GTCGTTCCCTGTGGTGTCTCAGAT
T16	GGTGAACGCTGTAGTTGGAATATA	GGTGAACGCTCAGATTTAAATATAAT
WK9	CCCGCAGTTAGATGCACCATT	CCGCAGTTAGATCAAGTGGC
P3	AACGGGCCAAAAACGGAGGT	AACGGGCCAACCGAG
TreB	CACTCCAGTTCTGCTCAAA	CACCTCCAAAACGAATATGG
AMs	CTTCCAAGGACCCCATCCT	CCCAACATCTCCGTCAGAAT
GPA	CCAAATACGCGGCCTTCT	AGTTTCTGGGCTCGGAGGA
AcPh	AGTTGTGGTTTAAAGCATAGG	ATTGTCCTTTTTCTTTAAAGTTTATTA
S3cI	CCACTCTCATGTCCCTTGAAC	GCCATGACATTTGGACAT
S3cII	TTCCATGATGTCCACTCTC	GGACAAATGTTTTCAGTGAATAAAT
GBSS1	CAAATAGCCACCCACACCAC	CTGTCAGATGTTCTTCTGATG
PP2	TTTGAATAGGTCCACTGCTT	CCATGCATCTCATTAGTCAA
PFruc	CCTTCTTCTTCGGGTGTCTCG	TGTTAAGTCCAGGGCAGAGG
Aglu	CCTCTGGAATCTTGCTATTTAGG	ATCCGCTAGATCACTGACAAA
LDS	CGAGGAGACAGACAGCATCA	GATGCAGCAGCCATGAGTT
BE2	GCCCCGAACATGATTCTA	GGCTTACCAGCCTTACTGT
Isa	CCTGTCTTGACGTCGCGTA	GCACGGTCTGATGTACGAGAG
SS1	TCTAGATTGCTACACGTGAGAGG	TCTCCACGATAACTTCCACC
SBE3	TCGGTCAATTCGGTTAGTCTCCTC	ACATCCTCTAGCATACTGGCGACTC
P3A	AATCCAACGCATCAAGGTGGC	ACAATGCCAAAACACCAGGAACCTCG
P4	TGAGCTTTACCTCCCCTCCTAACC	TCCACCTTCTCTCTCATCCAC
Pul3	GGGTTCGCTTTCACAACACAG	GTCACGACATAAGAGAAGCTGC
CBG	AGCTTCCCTAATGGCTTCGT	ATTTGCCAACTTTTGGATGG
HP	TGGAGGAGATGTACGTCGAG	GAAGTCGAGGTGGTCCATGA
P7	AGTTAAACAACCTCCCCTACTGC	GGGTAGGATAGGGGATAAGGAGC

changes from reference and alternate alleles, following the previous scoring method (Lestari *et al.* 2009). This study focused on formulation of regression equations using the binary data from a total 30 molecular markers, and the statistical means of physicochemical properties of 24 rice varieties. Thus, for the values obtained for measurements of AC, PC, and six RVA pasting properties (PV, HPV, CPV, BDV, SBV, and CTV) for each of the studied rice varieties, we simply calculated the range and mean. We searched for patterns of variation in the rice varieties by analyzing our genotyping data from DNA markers as well as the physicochemical property measurements, using Principal Component Analysis (PCA) in NTSYS (Exeter Software, Setauket, NY) package (Rohlf 1993). Then, multiple regression analysis was performed to establish any association between molecular markers and physiochemical traits, using SAS software (SAS 2002).

For statistical analysis, binary data of 30 markers along with mean values of the measurements for each trait for each rice variety, were formatted in

an Excel file and exported to SAS, then analyzed to determine  $R^2$  values that would indicate a significant association of combination of markers. Several outputs of primer combinations from the lowest to highest  $R^2$  were obtained. Then, selection of specified number of primers combinations for each trait was carried according to the highest  $R^2$  which showed a relatively small number of primers in the combination. By using multiple regression analysis (SAS 2002), the best model equation for prediction was identified using physicochemical properties/scores as dependent variables and the binary data from molecular markers as independent variables. Summary statistics, the parameter estimate with significant t value of each marker for every combination led to the creation of an equation:  $Y = a + bX_1 + cX_2 + \dots + X_n$  where Y= the calculated value of each trait which is predicted; a= intercept of total markers in the set; b, c, ..., n = parameter estimate value in the corresponding primer;  $X_1, X_2, \dots, X_n$  = binary data of each primer (zero, 0 or one, 1); plus (+) and minus (-) denote math symbols shown in parameter estimates.

## RESULTS

**Physicochemical Properties of Rice.** Since phenotypic characters contributed in formulation of model equations using molecular markers, 24 Indonesian indica varieties with defining food and cooking characteristics related to physicochemical properties were selected according to the available phenotypic description of each variety (Table 1). In addition to two landraces of different texture (Rojolele and Kalimutu), there were 22 Indonesian-bred varieties with a range of AC from low (<20%) to high (>25%) and distinctive texture ranging from fluffy to separate (Allidawati & Bambang 1989). Most of the varieties were types adapted for lowland cultivation, and for comparison we also included a tidal swamp variety (Batanghari) and two upland varieties (Jatiluhur and Cirata). Most varieties had cereh as typical indica type, and only one variety is hairy grain surface, javanica type (Rojolele) selected for its aromatic and good taste.

The measurement of physicochemical properties of our rice varieties (Table 3) differed somewhat in AC value, from that given by the description of the varieties but were relatively comparable. These varieties showed ranges of AC, PC, and RVA pasting-property values of indica rice. Of all of the physicochemical properties, only AC was positively skewed in the varieties. AC and PC in the varieties ranged from 16.63 (Sintanur) to 26.51% (Tukad Balian) and 5.53 (Jatiluhur) to 10.33% (Batang

Gadis) with averages of 21.38% and 8.84%, respectively. Pasting viscosity values including PV, HPV, BDV, CPV, SBV, and CTV were 219.64, 127.58, 92.06, 255.24, 35.6, and 127.66 RVU, with a range of 124.16-305.03, 91.00-169.10, 32.91-150.08, 142.08-323.31, -26.25-100.75, and 31.06-172.45, respectively. These selected physicochemical properties were very important to be examined in order to support association analysis with molecular data for the formulation of marker set.

**Variation of Rice Varieties.** Molecular analysis using 30 markers showed genetic variation among 24 rice varieties. Examples of clear banding patterns produced by some markers in some varieties are presented in Figure 1. GPA and P3A were the polymorphic markers (Figure 1A,C) which showed differentiation among certain varieties, while S3cI was monomorphic (Figure 1B). The variation among varieties is very important to set the basis for marker set formulation, thus principal component analyses (PCA) were generated based on physicochemical properties and genotyping data from total molecular markers (Figure 2). PCA showed distinctive relative positions for the varieties for both physicochemical properties and molecular characteristics. Regarding physicochemical properties, PCA revealed that the Sintanur variety had very low AC, PC, SBV, and CTV values, which were much lower than those of most other varieties, but similar to the tidal swamp variety Batanghari, characterized by separated-

Table 3. Values of physicochemical properties of 24 indica rice varieties

Variety	AC (%)	PC (%)	PV (RVU)	HPV (RVU)	BDV (RVU)	CPV (RVU)	SBV (RVU)	CTV (RVU)
Rojolele	22.53	9.73	245.00	139.53	105.47	271.08	26.08	131.55
Ciliwung	18.72	8.67	265.41	156.42	109.00	296.03	30.61	139.61
Cisokan	21.89	7.72	138.78	91.00	47.78	235.18	96.40	144.18
Cibodas	23.42	8.03	160.72	109.09	51.63	256.47	95.75	147.39
Jatiluhur	24.21	5.53	225.50	150.86	74.64	323.31	97.81	172.45
Kalimutu	23.04	6.17	198.33	114.75	83.58	234.55	36.22	119.80
Cirata	20.41	9.25	230.69	114.64	116.05	221.31	-9.39	106.67
Memberamo	19.00	10.22	245.19	147.31	97.89	266.00	20.81	118.69
Ciherang	18.31	9.71	233.35	119.33	114.02	230.14	-3.21	110.80
Sintanur	16.63	6.43	204.78	111.03	93.75	142.08	-62.70	31.06
Cimelati	18.16	10.23	245.67	135.89	109.78	251.28	5.61	115.39
Maros	22.88	9.73	170.05	93.97	76.08	224.78	54.72	130.80
Singkil	21.25	9.93	254.75	169.10	85.65	317.22	62.47	148.12
Batanghari	21.35	8.43	305.03	154.95	150.08	286.94	-18.08	132.00
Conde	21.48	9.07	207.55	119.83	87.72	258.08	50.53	138.25
Angke	21.30	10.14	202.25	113.92	88.33	229.25	27.00	115.33
Batang Gadis	20.87	10.33	222.64	141.88	80.76	284.02	61.38	142.14
Batang Piaman	24.11	8.01	255.50	144.69	110.81	280.39	24.89	135.69
Cigeulis	19.10	9.19	252.36	126.14	126.22	246.47	-5.89	120.34
Fatmawati	20.97	9.75	241.97	136.19	105.78	266.08	24.11	129.89
Konawe	20.66	9.74	215.06	116.33	98.72	241.11	26.06	124.78
Logawa	24.59	6.93	124.16	91.25	32.91	209.91	85.75	118.66
Pepe	21.76	9.78	202.72	131.41	71.31	258.03	55.31	126.62
Tukad Balian	26.51	9.57	224.03	132.47	91.56	296.11	72.08	163.64






Figure 1. Examples of banding pattern produced by primers GPA, S3cI, and P3A on some Indonesian indica rice varieties which were observed in this study. (A) Primer GPA (M. DNA-ladder of 100 bp, lane 1. Ciliwung, 2. Rojolele, 3. Jatiluhur, 4. Cibodas, 5. Kalimutu); (B) Primer S3cI (1. Ciliwung, 2. Rojolele, 3. Cisokan, 4. Cibodas, 5. Jatiluhur); (C) Primer P3A (lane 1. Ciliwung, 2. Rojolele, 3. Cisokan, 4. Cibodas, 5. Batanghari).

grain texture and low SBV. Some varieties such as Cibodas, Logawa, and Cisokan had physicochemical values that were notably more similar to each other than to other varieties. With regard to PCA values, the two landraces (Rojolele and Kalimutu) with different taste and texture, also were quite distinct in marker characterization each other and from the other improved varieties (Table 1, 3, Figure 2A,B). Marker genotyping was also able to identify the preferred taste varieties characterized by low AC and high PC. These varieties were Memberamo and Cimelati, clearly distinguished from the other varieties based on physicochemical properties (Figure 2B). Thus, these results suggest that there is a preliminary indication of interaction among 30 markers to differentiate rice with different genetic background and physicochemical properties.

#### Formulation of Molecular Marker Sets.

Association analysis between molecular markers and physicochemical properties was performed using the same method previously described to predict eating quality of japonica rice. In this analysis, only significant markers were included in the equations with parameter estimates used as regression coefficients for each marker (Table

4). The significant markers used as independent variables were able to generate equations to estimate each rice physicochemical property. The model equations that resulted from multiple regression analysis to predict physicochemical properties

Table 4. The parameter estimates and *t* value of markers used as significant coefficient

Trait	Primer	Parameter estimate	<i>t</i> value	
AC	E30	5.80 ± 0.74	7.79**	
	G4	2.94 ± 0.91	3.23**	
	G28	-3.93±0.44	-9.02**	
	AMs	6.10±0.61	10.07**	
	AcPh	4.94±0.65	7.64**	
	S3cI	-2.06±0.92	-2.24**	
	GBSS1	1.47±0.63	2.34**	
	PP2	10.29±1.78	5.77**	
	Pfruc	8.91±0.88	10.15**	
	Aglu	4.40±1.56	2.82**	
	BE3	0.84±0.71	1.18**	
	SS1	4.68±1.07	4.4**	
	Sbe3	-2.61±0.69	-3.76**	
	P3A	-3.89±1.43	-2.73**	
	P4	4.61±0.87	5.32**	
	Pul3	5.22±1.25	4.19**	
	CBG	4.72±0.61	7.69**	
	HP	-3.45±0.80	-4.32**	
	Isa	1.44±0.97	1.47**	
	Intercept		-0.92±3.18	-0.29**
	PC	F6	1.67±0.16	10.75**
		G4	3.42±0.21	16.5**
		S13	-1.16±0.15	-7.72**
		GPA	-2.49±0.36	-6.91**
		AcPh	1.05±0.16	6.5**
		S3cI	2.73±0.26	10.68**
		GBSS1	-2.75±0.17	-15.98**
		Pfruc	1.40±0.20	6.97**
		Aglu	-5.25±0.67	-7.81**
		LDS	11.39±0.60	18.9**
Sbe3		-1.91±0.28	-6.87**	
P3A		-4.05±0.31	-13.07**	
P4		3.76±0.26	14.25**	
Pul3		5.54±0.33	16.51**	
CBG		2.12±0.18	12.09**	
HP	-3.81±0.28	-13.82**		
Intercept		1.39±0.45	3.1**	
PV	A7	-107.51±12.33	-8.72**	
	E30	-112.93±7.37	-15.33**	
	G4	12.89±7.57	1.7**	
	G28	22.06±6.31	3.5**	
	S13	-35.92±4.71	-7.62**	
	WK9	33.36±7.48	4.46**	
	P3	12.25±5.42	2.26**	
	AMs	-71.00±7.81	-9.09**	
	GBSS1	-23.88±5.15	-4.63**	
	Pfruc	-91.77±5.74	-15.98**	
	LDS	33.06±13.97	2.37**	
	Isa	-235.88±15.41	-15.31**	
	SS1	-193.75±11.06	-17.52**	
	P4	-29.58±8.88	-3.33**	
	CBG	-37.49±6.74	-5.56**	
HP	-46.41±6.46	-7.18**		
Intercept		756.13±33.35	22.67**	

Table 4. Continued

Trait	Primer	Parameter estimate	<i>t</i> value	
CTV	A7	-122.61±8.56	-14.32**	
	F6	69.36±2.42	28.68**	
	G4	24.22±3.89	6.22**	
	S13	20.12±2.24	8.97**	
	WK9	31.33±3.30	9.48**	
	P3	24.91±2.09	11.91**	
	TreB	40.64±6.18	6.58**	
	AMs	19.47±2.15	9.05**	
	GPA	-85.74±5.04	-17**	
	AcPh	39.50±2.11	18.73**	
	SS1	-77.60±5.00	-15.53**	
	Sbe3	-6.27±3.19	-1.96**	
	P4	79.78±3.73	21.37**	
	Pul3	72.56±4.87	14.89**	
	CBG	-17.25±2.73	-6.32**	
	HP	-18.23±3.14	-5.8**	
	Intercept	85.36±8.92	9.57**	
	BDV	F6	-39.75±3.16	-12.58**
		G4	42.39±2.68	15.83**
G28		-6.77±2.56	-2.64**	
S13		-57.37±2.08	-27.64**	
T16		92.02±4.85	18.96**	
TreB		131.15±6.68	19.65**	
AcPh		32.97±2.28	14.47**	
S3cI		57.69±4.94	11.67**	
GBSS1		-46.05±2.61	-17.66**	
Pfruc		35.73±2.44	14.62**	
BE3		-10.15±2.47	-4.10**	
P3A		-97.79±3.84	-25.48**	
Pul3		28.32±3.76	7.53**	
CBG		30.24±2.50	12.09**	
HP		-56.94±3.49	-16.32**	
Intercept		126.39±8.19	15.43**	
SBV		F6	124.93±4.58	27.27**
		G4	-13.10±4.57	-2.87**
		G28	51.62±3.71	13.91**
	S13	58.96±3.32	17.73**	
	WK9	-24.41±6.00	-4.07**	
	P3	34.52±3.24	10.65**	
	AcPh	-13.29±4.30	-3.09**	
	Pfruc	-41.79±4.01	-10.41**	
	Aglu	57.76±9.21	6.27**	
	Isa	100.97±6.66	15.17**	
	Sbe3	-36.04±4.93	-7.30**	
	P3A	85.67±6.87	12.46**	
	P4	75.88±6.89	11.01**	
	Pul3	19.36±6.83	2.83**	
	CBG	-10.87±4.06	-2.67**	
	HP	26.56±5.56	4.78**	
	Intercept	-327.28±14.02	-23.34**	
	HPV	A7	-84.58±16.76	-5.05**
		E30	-57.44±5.27	-10.86**
G4		42.75±4.83	8.86**	
T16		52.03±18.49	2.81**	
AMs		-13.29±3.83	-3.47**	

Table 4. Continued

Trait	Primer	Parameter estimate	<i>t</i> value
HPV	GPA	-57.97±6.59	-8.80**
	AcPh	6.57±3.62	1.81**
	S3cI	-32.33±6.68	-4.84**
	GBSS1	-20.46±3.37	-6.07**
	PP2	51.80±8.75	5.92**
	Pfruc	-49.20±4.32	-11.38**
	SS1	-88.14±10.96	-8.04**
	P4	-29.41±6.30	-4.67**
	P7	11.97±7.00	1.71**
	Pul3	33.24±8.96	3.71**
	HP	-36.25±5.57	-6.51**
	Intercept	247.89±15.75	15.74**
CPV	A7	-109.18±15.68	-6.96**
	E30	-52.52±6.95	-7.56**
	F6	78.46±6.76	11.61**
	G4	65.81±8.77	7.50**
	G28	21.24±5.07	4.19**
	P3	27.09±5.66	4.79**
	TreB	170.20±18.16	9.37**
	GPA	-123.19±10.76	-11.45**
	AcPh	35.87±5.63	6.38**
	GBSS1	-45.98±5.11	-8.99**
	Pfruc	-43.68±6.25	-6.99**
	BE3	14.33±8.65	1.66**
SS1	-117.55±13.84	-8.49**	
Sbe3	-41.17±8.75	-4.70**	
P4	47.04±10.90	4.31**	
Pul3	87.20±11.16	7.81**	
CBG	10.51±6.09	1.73**	
HP	-66.78±8.54	-7.82**	
Intercept	240.06±29.16	8.23**	

\*\*significant at 1% level.

such as AC, PC, and six RVA pasting properties of indica rice are shown in Table 5. Highly significant resolution of each equation ( $R^2 = 0.99$ ) was obtained for prediction of AC, PC, PV, CPV, BDV, HPV, and CTV, and  $R^2 = 0.98$  for SBV. These results suggest that the physicochemical properties values of 24 varieties can be predicted reliably by these regression equations built with marker genotype data. The marker sets for the equations consisted of 15 markers for BDV, 16 markers for PC, PV, SBV, HPV, and CTV. CPV and AC could be estimated by a combination of 18 and 19 markers, respectively. Interestingly, the marker sets for the equations were dominated by 12 markers that we developed on the basis of eating quality-QTLs and candidate genes associated with the traits (TreB, GPA, AMs, AcPh, S3cI, GBSS1, PP2, PFruc, AGlu, CBG, HP, and LDS). Among markers included in the equations,




Figure 2. Principal component analysis (PCA) of 24 Indonesian-bred indica rice varieties based on physicochemical properties (A) and 30 molecular markers (B).

two markers (PFruc and HP) were shared for most traits but not for CTV. AcPh and CBG dominated most equations for evaluation of physicochemical properties excluding PV and HPV, respectively.

## DISCUSSION

High rice quality, attributable to preferred physicochemical properties, is necessary to meet high consumer demand for desired palatability and cooking quality of rice. Evaluation of physicochemical properties has long been conducted on japonica and indica rice by well understood and

commonly used means of measurement. However, conventional methods to determine for PC, AC, and major descriptors of pasting properties are not efficient. Molecular markers may provide an alternative approach for predicting rice starch characteristics. At present, several molecular marker sets have been established for prediction of japonica rice palatability (Ohtsubo *et al.* 2002, 2003; Lestari *et al.* 2009).

The wide range of physicochemical properties of 24 indica rice varieties used in this study, along with different genetic profiles, provides a good basis for association analysis of physicochemical



Table 5. Model equations containing significant molecular markers generated by multiple regressions analysis for prediction of physicochemical properties of Indonesian-bred indica rice

Trait*	Equation	R <sup>2</sup>
AC	AC= -0.92 + 5.80(E30) + 2.94(G4) -3.93(G28) + 6.10(AMs)+4.94(AcPh)-2.06(S3cI)+1.47(GBSS1)+10.29 (PP2)+8.91(Pfruc)+4.40(Aglu)+0.84(BE3)+4.68(SSI)-2.61(Sbe3)-3.89(P3A)+4.61(P4)+5.22(Pul3)+4.72 (CBG)-3.45(HP)+1.44(Isa)	0.99
PC	PC= 1.39+1.67(F6)+3.42(G4)-1.16(S13)-2.49(GPA)+1.05(AcPh)+2.73(S3cI)-2.75(GBSS1)+1.40(Pfruc)-5.25(Aglu)+11.39(LDS)-1.91(Sbe3)-4.05(P3A)+3.76(P4)+5.54(Pul3)+2.12(CBG)-3.81(HP)	0.99
PV	PV= 756.13-107.51(A7)-112.93(E30)+12.89(G4)+22.06(G28)-35.92(S13)+33.36(WK9)+12.25(P3)-71.0(AMs)-23.88(GBSS1)-91.77(Pfruc)+33.06(LDS)-235.88(Isa)-193.75(SSI)-29.58(P4)-37.49(CBG)-46.41(HP)	0.99
CTV	CTV= 85.36-122.61(A7)+69.36(F6)+24.22(G4)+20.12(S13)+31.33(WK9)+24.91-(P3)+40.64(TreB)+19.47(AMs)-85.74(GPA)+39.5(AcPh)-77.6(SSI)-6.27(Sbe3)+79.78(P4)+72.56(Pul3)-17.25(CBG)-18.23(HP)	0.99
BDV	BDV= 126.39-39.75(F6)+42.39(G4)-6.77(G28)-57.37(S13)+92.02(T16)+ 131.15(TreB)+ 32.97(AcPh)+57.69(S3cI)-46.05(GBSS1)+35.73(Pfruc)-10.15(BE3)-97.79(P3A)+28.32(Pul3)+30.24(CBG)-56.94(HP)	0.99
SBV	SBV=-327.28+124.93(F6)-13.10(G4)+51.62(G28)+58.96(S13)-24.41(WK9)+34.52(P3)-13.29(AcPh)-41.79(Pfruc)+57.76(Aglu)+100.97(Isa)-36.04(Sbe3)+85.67(P3A)+75.88(P4)+19.36(Pul3)-10.87(CBG)+26.56(HP)	0.99
HPV	HPV=247.89-84.58(A7)-57.44(E30)+42.75(G4)+52.03(T16)-13.29(AMs)-57.97(GPA)+6.57(AcPh)-32.33(S3cI)-20.46(GBSS1)+51.80(PP2)-49.20(Pfruc)-88.14(SSI)-29.41(P4)+11.97(P7)+33.23(Pul3)-36.25(HP)	0.98
CPV	CPV= 240.06-109.18(A7)-52.52(E30)+78.46(F6)+65.81(G4)+21.24(G28)+27.09(P3)+170.20(TreB)-123.19(GPA)+35.87(AcPh)-45.98(GBSS1)-43.68(Pfruc)+14.33(BE3)-117.55(SSI)-41.17(Sbe3)+47.04 (P4)+87.20(Pul3)+10.51(CBG)-66.78(HP)	0.99

\*AC: amylose content, PC: protein content, PV: peak viscosity, CTV: consistency viscosity, BDV: breakdown viscosity, SBV: setback viscosity, HPV: hot peak viscosity, CPV: cook peak viscosity.

properties with molecular data. The varieties used were diverse with regard to historical indica type (cereh and hairy grain surface), days to maturity, and suitability for cultivation in different habitats: either upland, lowland, or tidal swamp. Most of varieties had medium AC and PC, typical indica (cereh), and high yield potency (Table 1 & 3). The number of varieties analyzed genetically in this study is similar to the number used for indica rice taste estimation of 22 varieties which proved to be enough to establish marker sets (Lestari *et al.* 2009).

AC, PC and pasting properties are usually measured for routine analyses when screening germplasm, as part of the selection process in rice breeding. All physicochemical properties of starch differ by plant variety, each of which possesses a different genetic profile as well. AC and PC values seemed to relate to different patterns of rice grain quality. Most of the RVA pasting properties have been attributed to palatability, whereas AC levels tended to affect cooking quality (Allahgholipour *et al.* 2006; Lestari *et al.* 2009; Sun *et al.* 2011). In indica rice germplasm for large-grain varieties, pasting properties have been shown to relate to more or less gelatinous consistency (Allahgholipour *et al.* 2006; Lin *et al.* 2010). Differences in magnitude of the values for pasting properties provide additional characters for differentiating rice grain (Gravois & Webb 1997). Overall, AC and pasting properties are good indicators to measure rice cooking quality.

Thus, the selection of plant materials with diverse physicochemical properties was valuable in order to formulate market sets in regression analysis.

The high fidelity of molecular markers in contrast to nucleotide variation on the genome is useful for various genetic studies, and in particular for differentiating varieties in our research. Several molecular markers associated with physicochemical properties and overall rice eating and cooking qualities have been developed. These markers were derived from several genes that are associated with physicochemical properties of rice such as starch synthesizing genes (He *et al.* 2006; Bao *et al.* 2006a,b) and rice palatability (Ohtsubo *et al.* 2002, 2003). The markers used in our study were developed on the basis of genes linked to physicochemical properties-QTL, genes involved in starch synthesis in the rice pathway, and markers associated with rice physicochemical properties. Therefore, these markers are likely to hold potential for the development of PCR-based methods of prediction not only of rice palatability but also of the specific physicochemical properties of both indica and japonica rice.

A total of eight parameters, namely PC, AC, and six RVA pasting viscosities (PV, CPV, BDV, HPV, CTV, and SBV) were used for PCA of 24 indica varieties, and compared with molecular markers identified using PCA. This useful tool allowed for visualization of the differences among varieties, as

well as possible group affinities, and relationships among individuals (Li *et al.* 2011). The dot plots for PCA based on marker data and physicochemical trait data were distinguishable. The plot of DNA marker data tended to pool of most varieties on the basis of shared genetic background and characteristics, whereas the plot of physicochemical properties revealed more dispersed varieties. Markers were able to separate clearly two landraces (Rojolele and Kalimutu) and two improved varieties with preferred taste characteristics, from other improved varieties that could not be otherwise identified by physicochemical properties. This PCA indicates that DNA markers could be a valuable tool to complement differentiation of varieties based on physicochemical characters. A previous study evaluating japonica rice eating quality showed that examining the genetic relationship among varieties based only on data from a single marker may not be effective (Lestari *et al.* 2009), therefore, a set of multiple markers is needed for rice varieties differentiation.

A combination of markers is better than individual markers to estimate rice eating quality. Multiple regression analyses for association of a number of traits and molecular data have been performed (Tuyen & Prasad 2008; Fahlani *et al.* 2011) and have proved to be rapid and effective for predicting rice eating qualities using formulated markers (Ohtsubo *et al.* 2003; Lestari *et al.* 2009; Sun *et al.* 2011). The same approach may be applicable for evaluation of indica rice physicochemical properties.

The model equations formulated in this study included most of our previously developed markers based on eating quality-QTL and selected candidate genes in rice pathway. Our equations had high regression coefficients ranging from 0.98 to 0.99 (Table 5). The regression equations contained 15 to 19 significant markers, a feasible number for PCR amplification. These markers sets are comparable with marker sets containing 13 and 14 markers in the sets for estimation of japonica rice eating quality based on the Toyo taste meter, and sensory test, respectively (Lestari *et al.* 2009). In particular, the marker set based on Toyo taste meter after validation, can be used as a tool in the service of sample analysis for evaluation of Korean rice germplasm and populations. Clearly, even though these markers were developed from both japonica and indica rice genomes (Ohtsubo *et al.* 2002, 2003; Ohtsubo & Nakamura 2007; Lestari *et al.* 2009) they worked well on indica varieties.

A number of markers such as PFruc, HP, AcPh and CBG, were shared among equations. Some CAPS/dCAPS markers transformed into dominant SNAP markers (Lestari & Koh 2013) and included in the marker sets clearly may increase the reliability of the formulated marker sets. Since these regression equation results were part of a preliminary analysis, these marker sets may possibly be reformulated using a larger number of rice accessions. Further studies are also needed to confirm our findings and validate the formulated marker sets in the equations using more diverse indica rice varieties and breeding lines. PCR analysis requires much less sample material for testing, and allows the use of leaves rather than rice grains as a source of genomic DNA. This means testing can be conducted in early generations of population-crossing, which makes this tool even more promising. Ultimately, the marker sets in our study could be useful for assisting in the selection process to enhance breeding for indica rice varieties with improved eating and cooking qualities in Indonesia in the future.

## ACKNOWLEDGEMENT

This work was supported by a grant (No. 609001-5) from the Agricultural Research and Development Promotion Center for Agriculture and Forestry, Ministry of Agriculture and Forestry, Republic of Korea.

## REFERENCES

- Allahgholipour M, Ali AJ, Alinia F, Nagamine T, Kojima Y. 2006. Relationship between rice grain amylose and pasting properties for breeding better quality rice varieties. *Plant Breed* 125:357-262. <http://dx.doi.org/10.1111/j.1439-0523.2006.01252.x>
- Allidawati, Bambang K. 1989. Metode uji mutu beras dalam program pemuliaan padi. Dalam: Ismunadji M, Syam M, Yuswadi (Eds). *Padi Buku 2. Pusat Penelitian dan Pengembangan Tanaman Pangan*, Bogor. p 363-375.
- [AOAC] Association of Official Analytical Chemists. 1995. AOAC Official Methods of Analysis. 16th ed.; AOAC: Gaithersburg, MD. p 30.
- Bao JS, Corke H, Sun M. 2006a. Microsatellites, single nucleotide polymorphisms and a sequence tagged site in starch-synthesizing genes in relation to starch physicochemical properties in non-waxy rice (*Oryza sativa* L.). *Theor Appl Genet* 113:1185-1196. <http://dx.doi.org/10.1007/s00122-006-0394-z>
- Bao JS, Corke H, Sun M. 2006b. Nucleotide diversity in starch synthase IIa and validation of single nucleotide polymorphisms in relation to starch gelatinization temperature and other physicochemical properties in rice (*Oryza sativa* L.). *Theor Appl Genet* 113:1171-1183. <http://dx.doi.org/10.1007/s00122-006-0394-z>

- Bao JS, Jin L, Xiao P, Shen S, Sun M, Corke H. 2008. Starch physicochemical properties and their associations with microsatellite alleles of starch-synthesizing genes in a rice RIL population. *J Agric Food Chem* 56:1589-1594. <http://dx.doi.org/10.1021/jf073128+>
- Bao JS, Xia YW. 1999. Genetic control of the paste viscosity characteristics in indica rice (*Oryza sativa* L.). *Theor Appl Genet* 98:1120-1124. <http://dx.doi.org/10.1007/s001220051175>
- Fahliani RA, Khodambashi M, Houshmand S, Arzani A, Sorkheh K. 2011. Heritability for some agronomic characters of rice (*Oryza sativa* L.) and their linked microsatellites identification. *Turk J Agric For* 35:481-490.
- Gravoys KA, Webb BD. 1997. Inheritance of long grain rice amylograph viscosity characteristics. *Euphytica* 97:25-29. <http://dx.doi.org/10.1023/A:1003082930147>
- He Y, Han YP, Jiang L, Xu CW, Lu JF, Xu ML. 2006. Functional analysis of starch-synthesis genes in determining rice eating and cooking qualities. *Molecular Breeding* 18:277-290. <http://dx.doi.org/10.1007/s11032-006-5505-7>
- Izawa T. 2008. The process of rice domestication: a new model based on recent data. *Rice* 1:127-134. <http://dx.doi.org/10.1007/s12284-008-9014-7>
- Kuo BJ, Hong MC, Thseng FS. 2001. The relationship between the amylographic characteristics and eating quality of japonica rice in Taiwan. *Plant Prof Sci* 4:112-117. <http://dx.doi.org/10.1626/pps.4.112>
- Lestari P, Ham TH, Lee HH, Woo MO, Jiang WJ, Chu SH, Kwon SK, Ma KH, Lee JH, Cho YC, Koh HJ. 2009. PCR marker-based evaluation of eating quality of cooked rice (*Oryza sativa* L.). *J Agric Food Chem* 57:2754-2762. <http://dx.doi.org/10.1021/jf803804k>
- Lestari P, Koh HJ. 2013. Development of new CAPS/dCAPS and SNAP markers for rice eating quality. *Hayati J Biosci* 20:15-23. <http://dx.doi.org/10.4308/hjb.20.1.15>
- Li Z, Wan J, Xia J, Yano M. 2003. Mapping of quantitative trait loci controlling physicochemical properties of rice grains. *Breeding Sciences* 53:209. <http://dx.doi.org/10.1270/jsbbs.53.209>
- Li ZM, Zheng XM, Ge S. 2011. Genetic diversity and domestication history of African rice (*Oryza glaberrima*) as inferred from multiple gene sequences. *Theor Appl Genet* 123:21-31. <http://dx.doi.org/10.1007/s00122-011-1563-2>
- Lin Q, Liu ZH, Xiao H, Li L, Yu FX, Tian W. 2010. Studies on the pasting and rheology of rice starch with different protein residual. *IFIP AICT* 317:407-419.
- Mohapatra D, Bal S. 2011. Physical properties of indica rice in relation to some mechanical properties indicating grain characteristics. *Food and Bioprocess Technol* <http://dx.doi.org/10.1007/s11947-011-0539-1215>.
- Murray MG, Thompson WF. 1980. Rapid isolation of high molecular weight DNA. *Nucleic Acids Res* 8:4321-4325. <http://dx.doi.org/10.1093/nar/8.19.4321>
- Ohtsubo K, Nakamura S. 2007. Variety identification of rice (*Oryza sativa* L.) by Polymerase Chain Reaction method and its application to processed rice products. *J Agric Food Chem* 55:1501-1509. <http://dx.doi.org/10.1021/jf062737z>
- Ohtsubo K, Nakamura S, Imamura T. 2002. Development of the primer sets for identification of a rice variety, Koshihikari, by PCR. *Nippon Nogeikagaku Kaishi* 76:388-397. <http://dx.doi.org/10.1271/nogeikagaku1924.76.388>
- Ohtsubo K, Nakamura S, Okadome H. 2003. Investigation on estimation of rice palatability by DNA analysis (studies on estimation of rice palatability by DNA analysis part I). *Nippon Nogeikagaku Kaishi* 50:122-132.
- Patindol J, Guraya H, Champagne E, Chen MH, McClung A. 2010. Relationship of cooked-rice nutritionally important starch fractions with other physicochemical properties. *Starch/Stärke* 62:246-256. <http://dx.doi.org/10.1002/star.200900260>
- Perez CM, Juliano BO. 1978. Modification of the simplified amylose test for milled rice. *Starch/Stärke* 30:424-426. <http://dx.doi.org/10.1002/star.19780301206>
- Pooni HS, Kumar I, Khush GS. 1993. Genetical control of amylose content in selected crosses of indica rice. *Heredity* 70:269-280. <http://dx.doi.org/10.1038/hdy.1993.39>
- Ramesh M, Ali SZ, Bhattacharya KR. 1999. Structure of rice starch and its relation to cooked-rice texture. *Carbohydr Polym* 38:337-347. [http://dx.doi.org/10.1016/S0144-8617\(98\)00125-8](http://dx.doi.org/10.1016/S0144-8617(98)00125-8)
- Rohlf FJ. 1993. NTSYS-PC Numerical Taxonomy and Multi Variate Analysis System, version 1.8; Exeter Publishers: Setauket, NY.
- [SAS] Statistical Analysis Systems. 2002. SAS User's Guide, release 8.2; Statistical Analysis Systems Institute: Cary, NC.
- Sun MM, Abdula SE, Lee HJ, Cho YC, Han LZ, Koh HJ, Cho YG. 2011. Molecular aspect of good eating quality formation in japonica rice. *PLoS One* 6, e18385. <http://dx.doi.org/10.1371/journal.pone.0018385>
- Syamsir E, Hariyadi P, Fardiaz D, Andarwulan N, Kusnandar F. 2012. Pengaruh proses heat-moisture treatment (HMT) terhadap karakteristik fisikokimia pati. *J Teknol Industri Pangan XXIII*:100-106.
- Tian R, Jiang GH, Shen LH, Wang LQ, He YQ. 2005. Mapping quantitative trait loci underlying the cooking and eating quality of rice using a DH population. *Molecular Breeding* 15:117-124. <http://dx.doi.org/10.1007/s11032-004-3270-z>
- Tuyen DD, Prasad DT. 2008. Evaluating difference of yield trait among rice genotypes (*Oryza sativa* L.) under low moisture condition using candidate gene markers. *Omonrice* 16:23-33.
- Zhao WG, Chung JW, Kwon SW, Lee JH, Ma KH, Park YJ. 2012. Association analysis of physicochemical traits on eating quality in rice (*Oryza sativa* L.). *Euphytica* <http://dx.doi.org/10.1007/s10681-012-0820-z>.