

ECONOMIC IMPACT FROM PLASTIC DEBRIS ON SELAYAR ISLAND, SOUTH SULAWESI

DAMPAK EKONOMI DARI SAMPAH PLASTIK DI PULAU SELAYAR, SULAWESI SELATAN

Roni Hermawan^{1*}, Ario Damar², and Sigid Hariyadi²

¹Post-graduate Student of Coastal and Marine Resources Management, FPIK-IPB, Bogor

*E-mail: rooney_hermawan@yahoo.com

²Departement of Aquatic Resources Management, FPIK-IPB, Bogor

ABSTRAK

Sampah plastik dalam jumlah besar terdeposit di pesisir Pulau Selayar yang berhadapan langsung dengan Laut Jawa selama musim barat. Sampah plastik telah menimbulkan dampak sosial dan ekonomi bagi nelayan di Pulau Selayar. Penelitian ini bertujuan untuk mengkaji komposisi, kepadatan dan sebaran dari sampah plastik; dampak terhadap sosial dan ekonomi. Metode transek garis digunakan dalam penelitian ini untuk menentukan jumlah dan sebaran sampah plastik. Ukuran sampah plastik yang diamati adalah >2,5 cm dikategorikan sebagai sampah makro. Penelitian ini dilaksanakan pada Februari sampai Maret 2016. Dampak sampah plastik menurunkan pendapatan dari pariwisata, industri perikanan, mengganggu operasi penangkapan ikan, memerlukan pembersihan dan perbaikan pada alat langkap. Biaya yang dikeluarkan untuk perbaikan dan pembersihan kapal ikan sekitar 192,9 juta rupiah tiap tahun dan perbaikan alat tangkap 156,2 juta rupiah per tahun. Sampah plastik terdiri dari botol plastik, gelas plastik, tali dan jaring ikan, korek gas, keranjang plastik, pelampung, kemasan plastik, sikat gigi dan alat suntik. Rata-rata sampah plastik adalah $9,5 \pm 2,7$ item/m² dan berat sekitar $229,2 \pm 109,9$ g/m².

Kata kunci: dampak, manajemen, Pulau Selayar, sampah plastik, sosial-ekonomi

ABSTRACT

Enormous plastic debris has been stranded on Selayar island shore during the west monsoon due to its position which is directly faced Java sea. Plastic debris had impacted social and economy of Selayar fishermen. The research was aimed to determine the composition, density, and distribution of the plastic debris; it impacts to social and economy. Line transect was used as the research method to quantify the numbers and width of the plastic debris spread. The size of plastic debris that was researched about >2,5 cm which categorized as macro litters. The research was conducted in February to March 2016. The impacts of plastic debris were reduce income from tourism, fishing industry, disrupting fishing operations, requiring clean-up, and repairing vessel. Direct costs for repair and clean-up fishing vessels was about 192.9 million rupiahs per year and repair fishing gears was about 156.2 million rupiahs per year. The plastic debris consists of plastic bottle, plastic cup, rope and fish net, gas matches, plastic box, buoy, food packaging, toothbrush and syringe. The average of plastic debris about 9.5 ± 2.7 item/m² and weight about 229.2 ± 109.9 g/m².

Keyword: impact, management, Selayar Island, plastic debris, socio-economy

I. INTRODUCTION

Plastic debris was increased as growing of economy and plastic for support human life as consumptions or other activities. Plastic products increase significant and domination of every market products

since 1930s to 1940s. World's plastic production reach 288 metric tonnes in 2012 or increase 620% from 1975 (Jambeck *et al.*, 2015). Plastic industries create new products to follow market trends and showing no signs of slowing (Moore, 2015). Plastic debris came from industry, domestic litter, shipping

vessel, boatship and another land-based source that drive by river or drainasse to the sea (Derraik, 2002 ; UNEP, 2005; Williams and Simmons, 1997).

Plastic debris has been deposit into ecosystem and degradation to be micro-plastic and become more dangerous. In the environmental, polymer-based spread everywhere and the impacts of polymer are yet fully understood (Lambert, 2013). The impacts of plastic debris to fishing industry are disturbing fishing vessel, ship propeller, ghost fishing from broken or abandoned fishing gear and repairing fishing vessel that requiring cost (Lee, 2014).

Coastal are a vulnerable area from marine waste, some of the ecosystems depend on coastal and used by many stakeholders (Lindgren, 2011). Plastic debris drive by surface currents, wind-driven mixing, down welling and/or currents act to carry plastic particles (Choy and Drazen, 2013),

then end up on the beach and make many environmental problems. West coast of Selayar island is directly facing open sea, on west monsoon (January to April) marine debris drive by surface current from Makassar strait and Java sea to Selayar island west coast (Balitbang KP, 2016; ECMWF, 2016; Wyrcki, 1961). Few researchs about marine plastic debris on Indonesia (Sherman and Seville, 2016; Uneputty and Evans, 1997; Uneputty and Evans, 1997b; Wala-langi, 2007; Willoughby, 1997).

Plastic litter on Selayar island west coast are unmanaged, when east monsoon (April to September) plastic litters debris mostly drive back by tide and surface current, then make another problem to other islands. The research was aimed to determine the types, quantities, and distribution of the plastic debris and its impacts to local socio-economic.




Figure 1. Point site research map.

II. METHODS

2.1. Time and Location of Research

The research was conducted on Selayar Island west coast in February to March 2016, it was the peak of west monsoon to east monsoon transition. The plastic litters was collected from 7 coastline sites depend on coastal landforms where plastic debris was stranded (Figure 1).

2.2. Methods and Analysis

Data of plastic debris were collected by line transect method to determine its numbers and spreads. Transects were 5 m width and length depend on coastline width. Sample sizes were macrodebris (> 2,5 cm), samples sorted by type for identification, measured, weighed (g) and measured the location plot site (m²). The plot point of observation was sorted to clean and marked, each station consists of three plots (Lippiatt *et al.*, 2013 and UNEP, 2009). Social economy impact observed by purposive sampling with 110 locals community, non-government organization, and local government.

2.2.1. Plastic Debris Density (Modified from Lippiatt *et al.*, 2013)

Data of plastic debris consist of numbers of debris (item), weight (g) and location area (m²).

$$C = \frac{n}{(w \times l)} \dots\dots\dots(1)$$

Where: C = Plastic debris density (item/m²); n = Number of debris (item); w = Transect width (m); l = Transect length (m).

2.2.1. Loss of Vessel (Modified from Takehama, 1990)

Datas of vessel were taken from vessel that operating and anchored on Selayar west coast, operated at less than 300 nautical miles or around of Selayar islands and South Sulawesi.

$$A = ((a \cdot b) \cdot N_1) + ((a \cdot c) \cdot N_2) \dots\dots\dots(2)$$

Where: A = Loss of vessel (unit); a = Σ vessel on west coast (unit); b = Frequency of; entangled of vessel propeller (unit); c = Frequency of others repairing/cleaning (unit); N₁ = Cost of propeller repairing (rupiah); N₂ = Cost of others repairing/cleaning (rupiah).

2.2.2. Loss of Fishing Gear (Modified from Takehama, 1990)

Fishing gear data were taken from fixed fishing gear such as set net, fixed lift net, boat lift net and fish traps that operated on Selayar west coast.

$$B = ((p \cdot q) \cdot R) \dots\dots\dots(3)$$

Where: B = Loss of fishing gear (rupiah); p = Σ fixed fishing gear on west coast (unit); q = Frequency of extra cleaning, fixing net, rame, buoy, anchor, etc (unit); R = Cost of gear repairing and cleaning (rupiah).

III. RESULT AND DISCUSSION

3.1. Types, Quantities and Distribution of Plastic Debris

Plastic litters consist of plastic bottle, plastic cup, rope and broken fish net, gas matches, plastic box, bouy, wrapper, tooth brush, and syringe (Figure 2 and 3). Most of the plastic litters are plastic for food or daily-need packaging, cups and bottle plastics are dominant. On site 4, total density was about 6 item/m² for plastic cup, site 7 plastic bottle was about 5 item/m², describe on Table 1.

On Table 3 shown weight (g/m²) of plastic litters, site 1 is highest with weight about 129 g/m² for rope and fish-net and 128,9 g/m² for plastic bottle. Rope and fish-net were from abandoned or parts of fishing gears. Based on the calculation results and total length of coastline per site (Figure 1), the total amount was about 7.310.173 item with average 9.5 ± 2.7 item/m² and total weight was about 241.759 kg with average 229.2 ± 109.9 g/m².

Table 1. Total of plastic debris per site (item/m²).

Type	Site						
	1	2	3	4	5	6	7
Plastic drink bottle	4.9	3.0	2.4	2.8	4.0	3.9	5.0
Plastic cup	3.7	4.7	5.5	3.6	6.0	4.3	3.2
Rope & fish-net	0.3	0.3	0.4	0.2	0.6	0.3	0.7
Gas matches	0.9	0.2	0.4	0.3	0.7	0.3	0.4
Basket	0.0	0.0	0.0	0.0	0.1	0.0	0.0
Bouy	0.04	0.02	0.00	0.03	0.05	0.03	0.03
Plastic packaging	0.7	0.3	0.4	0.4	0.4	0.2	0.3
Toothbrush	0.1	0.0	0.0	0.0	0.0	0.0	0.0
Plastic spoon	0.1	0.0	0.0	0.0	0.0	0.0	0.0
Syringe	0.1	0.0	0.0	0.0	0.0	0.0	0.0

Table 2. Weight of Plastic Debris per site (g/m²).

Type	Site						
	1	2	3	4	5	6	7
Plastic drink bottle	128.9	71.6	70.5	52.1	114.3	83.0	100.2
Plastic cup	36.6	52.3	32.7	32.7	62.8	44.3	51.6
Rope & fish-net	129.0	101.9	18.6	18.6	24.9	36.2	72.7
Gas matches	13.4	2.2	3.5	3.5	4.0	2.4	4.0
Basket	25.8	30.4	26.3	26.3	8.8	3.5	11.5
Bouy	18.0	6.4	0	0	16.0	6.2	14.3
Plastic packaging	11.7	2.0	2.7	2.7	3.1	7.8	8.1
Toothbrush	1.2	0	0	0	0.0	0	0
Plastic spoon	0.3	0.2	0	0	0.0	1.4	0
Syringe	0.4	0	0	0	0.0	0	0



Figure 2. Plastic debris stranded on the beach.



Figure 3. Broken fish-net abandoned on the beach.




Figure 4. Map of plastic debris spreads.

The spread of plastic debris along the west coast of Selayar island and deposited on the beach, west coast has a length of 108,59 km, plastic litter covered 70,78 km or 65,18% of coastline. Based on observations

and calculations at site 1 of 10 type of plastic debris, which was dominated plastic bottle (Figure 4b) and a highest average weight of plastic debris at site 1 was about 300-400 g/m^2 (Figure 4a).

3.2. Plastic Debris Impact on Social

Social appraisals for plastic debris quite varied, it is because some locals found that the deposited debris is a natural phenomenon that commonly occurs annually. To know the community assessment conducted by interviewing randomly at each site point of observation and categorized by Likert. Based on 7 observation site points with 110 respondents were mostly local who settled and lived on the west coast of Selayar island, presented in Figure 5. Aesthetics beach interrupted by marine debris were piled up, the plastic debris is also disturbing tourism activities such as swimming or enjoy the beach view. Local activities were interrupted by plastic debris as beach covered by plastic and marine debris, which is access to the shore and fishing activities. Fishing gears often disrupted and damaged by plastic debris, thus reducing catches. Marine litter provides the aesthetic impact on the coastal environment and the economic impact of various industries that depend on coastal environmental and sea (Bergmann, 2015).




Figure 5. Social impact of plastic debris.

Some type of are collected, utilized and sold by locals. Thick and heavy plastics are valuable, while plastics lightweight plastics such as mineral water bottle or plastic cups has no economic value and unmanaged while there were in large number, its can be potential benefit if it managed properly. Plastic waste collectors

only receive a thick plastic due to the cost of transport from Makassar to Selayar island. The collectors in Selayar island also brought commodity crops to Makassar for efficient transport costs and improve benefits. The collector bought plastic debris only as a side benefit.

3.3. Plastic Debris to Economy

The Majority of fishermen on Selayar island is a side job, most of the population relying on the farming as the main livelihood (BPS, 2015). Many fishermen come from small island around the Selayar island such as Tinabo, Pasimasunggu, Taka Bonerate and other islands. On the Selayar island boats and ships activities are very high because Selayar is the capital district of Selayar islands, thus becoming berths, fishing spot, transporting, docking and transaction from Makassar or other islands.

Floating plastic debris disturbing even damaging part of ship as propeller of ship, engine cooling system and other damage that requires repair and maintenance cost higher. Fishing gear also affected and require high costs. Based on calculation of loss cost of boat and fishing gear by plastic debris, presented on Table 3 and Table 4.

Losses of cost by boats and fishing gears (Table 3 and Table 4) derived from the number of vessels and fishing gears were operating on the west coast of Selayar island (>300 nautical miles) and information from local fishermen, shipowners, and ship docking worker.

Number of frequency entangled of vessel propeller and reaping costs from surveyed vessel that anchored on west coast of Selayar during west monsoon, information of entangled propeller was asked to helmsman, vessel crews and mechanic crews. Informations about loss of fixed fishing gear such as frequency of cleaning, fixing net, fixing frame, replacing bouy, anchor, rope, canopy and other parts were asked to gear owner, gear crews and other locals that worked on fixed fishing gear.

Table 3. Table of impacted fishing vessel by plastic debris.

Loss of Fishing-Vessel by Plastic Debris			
District	Small (<10 GT)	Moderate (10-30 GT)	Large (>30 GT)
Bontosikuyu (unit)	44	49	5
Bontoharu (unit)	59	77	9
Benteng (unit)	39	62	9
Bontomanai (unit)	15	45	10
Bontomatene (unit)	30	55	10
TOTAL	187	288	43
Amounts of vessel on west coast (unit)	47	130	37
Frequency of entangled of vessel propeller (unit)	0.05	0.09	0.25
Frequency of repairing (unit)	0.60	0.45	0.25
Cost of propeller repairing (Rp)	500.000	2.000.000	5.000.000
Cost of other repairing (Rp)	500.000	1.000.000	5.000.000
Numbers of entangled of vessel propeller (unit)	2	13	9
Numbers of others repairing/cleaning (unit)	28	58	9
Total cost propeller repairing (Rp)	1.168.750	25.920.000	46.762.500
Total cost for others repairing/cleaning (Rp)	14.025.000	58.320.000	46.762.500
Total (Rp)	1.5193.750	84.240.000	93.525.000
Total Loss Rupiah (Rp)	192.958.750		

Table 4. Table of impacted fishing gear by plastic debris.

Loss of Fishing Gear				
District	Fixed Lift-net (<i>Bagan</i>)	Boat Lift-net (<i>Bagan Apung</i>)	Set-Net (<i>Sero</i>)	Fish Traps
Bontosikuyu (unit)	5	5	3	12
Bontoharu (unit)	10	45	15	35
Benteng (unit)	3	5	-	-
Bontomanai (unit)	3	5	6	-
Bontomatene (unit)	0	0	25	-
Total fishing gear (unit)	21	60	49	47
Amounts of fishing gear on west coast (unit)	13	36	39	28
Impact and extra cleaning, net, frame, buoy, anchor, etc (unit)	45%	30%	80%	45%
Number of affected by plastic debris (unit)	6	11	31	13
Cost of repairing, fixing, cleaning (1 to 3 million rupiah)	17.010.000	32.400.000	94.080.000	12.690.000
Total Loss (rupiah)	156.180.000			

The total loss of the vessels were about 193 million IDR and fishing gears were about 156 million IDR each year. Nets of fishing gear often damaged by plastic debris such as stuck on the set-net that fixed installed in tidal areas, so its very easy affected by sharp or large number of plastic debris. Set net also easy to uprooted because its used wood-stick frame and require high cost for repairing.

Several studies of loss due to plastic debris from fishing vessels or fishing gears, such as the British ship losses 1.245 to 3.283 million US dollars in 2008 (Lee, 2014), in 2008, 286 rescues of vessels with fouled propellers in United Kingdom waters were carried out at a cost of between €830.000 and €2.189.000 (Watskins, 2015). Japan's loss 18,4 million US dollars for the ship in 1985 (Takehama, 1990). British losses for the fishing gear 12.000 to 67.500 US dollars in 2012 (Lee, 2014).

Plastic debris impacted all coastal activities such as marine organism, ecological processes, aesthetics problem, safety and health, decreasing number of tourism, ghost fishing, disturbing sea transportation. During west peroid there were no tourism activities on Selayar west coast because of marine litter. Abandoned, Lost or otherwise Discraded Fishing Gear/ ALDFG, fishing gear becomes ALDFG when the fisher loses all operational control of the equipment (Smolowitz *et al.*, 1978). As control over fishing gear is lost, the selectivity and efficiency of the gear for the original target species may be altered, overall ghost fishing catches are probably very low compared to controlled fishing (Brown *et al.*, 2005).

IV. CONCLUSION

Plastic debris on Selayar island spread along 70.78 km or 65.14% of west coast, dominated by plastic wrapper from daily consumption. Plastic debris density average 9.5 ± 2.7 item/m² and weight was about 229.2 ± 109.9 g/m². Socio-economic

impact to local activities, marine tourism, sea-transportation, ghost fishing and economic loss of fisheries industry especially for fishermen about 349 million IDR each year.

ACKNOWLEDGEMENT

The authors are grateful to all colleague for participating in this research and contributed to collecting Marine Litter survey data, such as crew of Takabonerate National Park, The government of Selayar Islands and Marine Coastal Management of IPB students.

REFERENCES

- Badan Penelitian dan Pengembangan Sumber Daya Laut dan Pesisir (Balitbang KP). 2016. Data Perairan Indonesia, <http://p3sdlp.litbang.kkp.go.id/index.php/data/perairan-indonesia>. [Retrieved on February 2016].
- Badan Pusat Statistik (BPS). 2015. Statistik daerah kabupaten Kepulauan Selayar 2015. Badan Pusat Statistik Kabupaten Kepulauan Selayar. Makassar. 40hlm.
- Bergmann, M., L. Gutow, and M. Klages. 2015. Marine anthropogenic litter. Springer international publishing AG Switzerland. Springer Science. Business Media. Gothenburg. 447p.
- Brown, J., G. Macfadyen, T. Huntington, J. Magnus, and J. Tumilty, 2005. Ghost fishing by lost fishing gear. Final Report to DG Fisheries and Maritime Affairs of the European Commission. Fish/2004/20. Institute for European Environmental Policy/ Poseidon Aquatic Resource Mana-gement Ltd joint report. www.ieep.org.uk/assets/243/ghostfishing.pdf. [Retrieved on October 2016].
- Choy, C. and D.J. razen. 2013. Plastic for dinner? Observations of frequent debris ingestion by pelagic predatory fishes from the central North Pacific.

- Marine Ecology Progress Series*, 485:155–163.doi:10.3354/meps1032.
- Derraik, J.G.B. 2002. The pollution of the marine environment by plastic debris: a review. *Marine Pollution Bulletin*, 44:842-852.
- European Centre for Medium-Range Weather Forecasts (ECMWF). 2015. Surface current data, <http://www.ecmwf.int/>. [Retrieved on February 2016].
- Jambeck, J.R., R. Geyer, C. Wilcox, T.R. Siegler, M. Perryman, A. Andrady, R. Narayan, and K.L. Law. 2015. Plastic Waste Inputs from Land into The Ocean. *Marine Pollution Science*, 347:768-771.
- Lambert, S. 2013. Environmental risk of polymers and their degradation Products. Dissertation. Environment Department. University of York, Canada. 198p.
- Lee, J. 2014. Economic valuation of marine litter and microplastic pollution in the marine environment: An initial assessment of the case of the United Kingdom. www.eftec.co.uk/lee-paper/download. [Retrieved on October 2016].
- Lindgren, D. 2011. Form and Function of coastal Areas. Dissertation. Upsalla University. Sweden. 52p.
- Lippiatt, S., S. Opfer, and C. Arthur. 2013. Marine Debris Monitoring and Assessment. NOAA Technical Memorandum NOS-ORandR-462005, http://marinedebris.noaa.gov/sites/default/files/Lippiatt_et_al_2013.pdf. [Retrieved on October 2016].
- Moore, C. 2015. How much plastic is in the ocean? You tell me. *Marine Pollution Bulletin*, 92:1–3.
- Sherman, P. and E. Seville. 2016. Modeling marine surface microplastic transport to assess optimal removal locations. *IOP Publishing. Environ. Res. Lett.* 11-19pp. DOI:10.1088/1748-326/11/1/014006.
- Smolowitz, R.J. Corps, L.N. and Center, N.F. 1978. Lobster, *Homarus americanus*, trap design and ghost fishing. *Mar. Fish. Rev.* 40(5):2–8.
- Takehama, S. 1990. Estimation of damages to fishing vessels caused by Marine Debris, Based on Insurance Statistics. *Fishing Ground Environment Conservation Division. Fisheries Agency Japan.* 792-808p.
- Uneputty, P.A. and S.M. Evans. 1997. Accumulation of Beach Litter on Islands of the Pulau Seribu Archipelago, Indonesia. *Marine Pollution Bulletin*, 34(8):652-655.
- Uneputty, P. and S.M. Evans. 1997b. The Impact of Plastic Debris on the Biota of Tidal Flats in Ambon Bay (Eastern Indonesia). *Marine Environmental Research*, 44(3):233-242
- United Nations Environmental Programme (UNEP). 2005. Marine litter: An analytical overview. http://www.unep.org/regionalseas/marinelitter/publications/docs/anl_oview.pdf. [Retrieved on February 2016].
- United Nations Environmental Programme (UNEP). 2009. Guidelines on survey and monitoring of marine litter. Regional Seas Reports and Studies No. 186 IOC Technical Series No. 83. www.unep.org/PDF/UNEP_2009_A_NNUAL_REPORT.pdf. [Retrieved on February 2016].
- Walalangi, J. 2012. Analisis komposisi sampah organik dan anorganik serta dampak terhadap lingkungan pesisir Kota Palu Sulawesi Tengah. Thesis. Bogor Agricultural University. 112p.
- Watkins, E., P.T. Brink, S. Withana, K. Mutafoglu, J.P. Schweitzer, D. Russi, and M. Kettunen. 2015. Marine litter: Socio-Economic Study. Scoping report. Institute for European Environmental Policy. London, Brussels. 365-394p.
- Williams, A.T. and S.L. Simmons. 1997. Estuarine litter at the river/beach

- interface in the Bristol Channel, United Kingdom. *J. of Coastal Research* 13:1159–1165.
- Willoughby, N.G., H. Sangkoyo, and B.O. Lakaseru. 1997. Beach litter: an increasing and changing problem for Indonesia. *Elsevier. Marine Pollution Bulletin*, 34:469–478.
- Wyrski. K. 1961. Scientific results of marine Investigations of the South China Sea and the Gulf of Thailand. 2nd ed. Pergamon Press Ltd. California. 195p.
- Diterima : 16 Maret 2017*
Direview : 22 Februari 2017
Disetujui : 20 Mei 2017