

SPATIAL AND TEMPORAL VARIATION OF ZOOPLANKTON COMPOSITION NEAR WHALE SHARK SIGHTINGS IN PROBOLINGGO OF EAST JAVA, INDONESIA

Nurlita Putri Anggraini^{1,4*}, Majariana Krisanti³, Hawis Madduppa²,
Mahardika Rizqi Himawan⁵

¹Study Program of Marine Science, Garduate School, IPB University, Bogor 16680, Indonesia

²Department of Marine Science and Technology, Faculty of Fisheries and Marine Science,
IPB University, Bogor 16680, Indonesia

³Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science,
IPB University, Bogor 16680, Indonesia

⁴Department of Marine Science, Faculty of Marine Science and Fisheries, University of Syiah Kuala,
Aceh, 23111, Indonesia

⁵Department of Marine Science, Faculty of Agriculture, Mataram University, Mataram, 83125,
Indonesia

*Corresponding author: nurlitaputrianggraini@gmail.com

(Received 21-06-2021; Revised 05-10-2023; Accepted 10-03-2024)

ABSTRACT

Whale shark occurrence in Probolinggo differs from other Indonesian locales, suggesting a link to zooplankton availability. Zooplankton composition and whale shark emergence are the focus of this study. From December 2017 to November 2018, six observation points were made each month. A plankton net filters and lugol preserves water. Olympus CX23 microscope observations were repeated twice. The spatial analysis revealed varying whale shark numbers at each station (Chi-square test, $X^2 = 1418.6$, $P < 0.05$), with six sharks observed at station PR_5. Zooplankton numbers were similar at each location. Temporal analysis revealed significant differences in whale shark appearance each month (Chi-square test, $X^2 = 81.04$, $P < 0.05$), with March and November having the highest appearance among the three individuals. The amount of zooplankton varied (Chi-square test, $X^2 = 148.61$, $P < 0.05$), with the highest abundance in April and March. Whale shark appearance and zooplankton composition were not correlated ($r = 0.01$, $P < 0.05$) both geographically and temporally. Whale sharks were linked to zooplankton kinds. Results indicate whale sharks are particularly interested in *Acartia* sp. ($r = 0.3$, $P < 0.05$). This suggests that whale sharks' appearance is determined by their demand for food, not zooplankton availability.

Keywords: aggregation, feeding habit, habitat use, *Rhincodon typus*

Variasi Spasial dan Temporal Komposisi Zooplankton di Wilayah Kemunculan Hiu Paus, Probolinggo, Jawa Timur, Indonesia

ABSTRAK

Pola kemunculan hiu paus di Probolinggo berbeda dengan lokasi lain di Indonesia, diduga kemunculannya memiliki hubungan dengan ketersediaan zooplankton. Tujuan penelitian ini adalah untuk mengkaji komposisi spasial dan temporal zooplankton serta mengaitkan dengan kemunculan hiu paus. Penelitian dimulai Bulan Desember 2017 - November 2018 dan terdapat enam titik pengamatan tiap bulannya. Air disaring menggunakan plankton net dan diawetkan menggunakan lugol. Pengamatan menggunakan mikroskop Olympus CX23 dengan dua kali ulangan. Hasil analisis spasial menunjukkan adanya variasi terhadap keberadaan hiu paus di setiap stasiun (Chi-square test, $X^2 = 1418.6$, $P < 0.05$) dengan kemunculan tertinggi diamati di stasiun PR_5 sebanyak enam individu. Namun, jumlah zooplankton tidak menunjukkan perbedaan yang signifikan di setiap stasiunnya. Berdasarkan analisis temporal, kemunculan hiu paus berbeda signifikan setiap bulannya (Chi-square

test, $X^2 = 81.04$, $P < 0.05$), dengan bulan Maret dan November menunjukkan kemunculan tertinggi, tiga individu. Terdapat variasi kelimpahan dalam jumlah zooplankton (Chi-square test, $X^2 = 148.61$, $P < 0.05$), dengan kelimpahan terbesar terjadi pada bulan April dan Maret. Baik secara spasial maupun temporal, korelasi kemunculan hiu paus terhadap komposisi zooplankton tidak berhubungan ($r = 0.01$, $P < 0.05$). Selain itu, korelasi antara hiu paus terhadap jenis zooplankton yang ditemukan juga dilakukan, hasilnya menunjukkan bahwa *Acartia* sp. terindikasi menjadi jenis target hiu paus tersebut ($r = 0.3$, $P < 0.05$), dimana kemunculan hiu paus tidak dipengaruhi oleh jumlah zooplankton yang tersedia, namun berdasarkan kebutuhan kalorinya.

Kata kunci: agregasi, kebiasaan makan, penggunaan habitat, *Rhincodon typus*

INTRODUCTION

The whale shark (*Rhincodon typus* Smith, 1828) exhibits a significant migratory pattern that is strongly influenced by the availability of food sources (Yap-dejeto *et al.*, 2013). Whale sharks are zooplankton-feeding organisms, as confirmed by fecal observations and abdominal exams (Nelson and Eckert, 2007; Boldrocchi *et al.*, 2020). Whale sharks consume many kilograms of zooplankton on a daily basis, as recorded by Heyman *et al.* (2001), Hacohe-Domene *et al.* (2006), Nelson and Eckert (2007), Hernandez-Nava and Alvarez-Borrego (2013), and Boldrocchi *et al.* (2020). The zooplankton species that whale sharks prey on include Copepod, crab larvae, fish eggs, fish larvae, Chaetognath, and others (Heyman *et al.*, 2001; Rowat and Engelhardt, 2007; Meekan *et al.*, 2009; Boldrocchi *et al.*, 2018). These findings suggest that the patterns of whale shark aggregation varied throughout each observation region. The groups of copepod, Chaetognatha, and Ctenophore have an impact on whale sharks at the observation location in Djibouti, North Africa (Boldrocchi *et al.*, 2020). The tintinnid group is the predominant species in the waters of the Philippines, which are also home to a significant population of whale sharks (Yap-dejeto *et al.*, 2013). Unlike the waters of the Ningaloo Reef, the whale shark shares its habitat with herring fish, which feed on the copepod found in these regions (Wilson, 2002). Nevertheless, the fact that the whale shark is an ophthalmological animal does not imply a specific preference for its diet. Its primary objective is not only

to devour huge amounts of food, but also to meet its energy requirements (Pierce and Norman, 2016).

Whale sharks inhabit several locations in Indonesia, including Cendrawasih Bay, Papua; Talisayan, Kalimantan; Botuborani, Gorontalo; and Bentar Beach, Probolinggo. In some locations, such as Cendrawasih Bay, Talisayan, and Botuborani, where local fishermen provide their catch, whale sharks no longer appear seasonally but permanently due to the presence of lifting nets (Himawan *et al.*, 2015). Even though there are lifting nets in Probolinggo, fishermen do not give their catch to whale sharks, so whale sharks appear seasonally in these waters (Kamal *et al.*, 2016). Researchers have conducted several studies on whale sharks in Probolinggo, including Kamal *et al.* (2016), Himawan (2017), Syah *et al.* (2018), and Kamal *et al.* (2020); however, only Kamal *et al.* (2016) and (2020) conducted research on zooplankton abundance at this location. Due to its opportunistic nature, the composition of zooplankton, or its preference for it, determines its appearance, causing it to change seasonally. Investigations into the spatial and temporal appearance of whale sharks in Probolinggo are necessary to complete and answer the question of their seasonal appearance. The aim of this study was to investigate a spatial and temporal of zooplankton composition in Bentar Beach, Probolinggo in six station for twelve months and then linking these findings to the presence of whale sharks. This research can provide additional data on whale sharks in Probolinggo and serve as the foundation for conservation efforts in this area.

RESEARCH METHODS

Study Site

The sampling stations were located in the tourist area of Bentar Beach, Probolinggo, which is part of the Madura Strait. We collected zooplankton samples at six stations (Figure 1) between December 2017 and November 2018. The station points selected were based on previous research (Kamal *et al.*, 2016; Himawan, 2017; Syah *et al.*, 2018; Kamal *et al.*, 2020) and reports from local people. We conducted the observation every month during daylight hours (06.00–14.00). The Laboratory of Marine Biodiversity and Biosystematics, Department of Marine Science and Technology, Faculty of Fisheries and Marine Sciences, Bogor Agricultural University conducted the observations.

size of 150 μm , a 50 cm diameter mouth conical net, and put into 100 ml bottle sampling. Lugol's solution was used to preserve the sample for taxonomical investigation. We have successfully identified the organism, conducted a thorough count, and calculated the density by expressing the number of individuals per cubic meter. For observations, we utilized an Olympus CX23 microscope set at a magnification of 10x. To collect the sample water, we obtained 1 ml and placed it in a Sedgewick-Rafter (SRC). Morphological observations were performed with identification books authored by Catellani and Edward (2017). We performed zooplankton identification at the genus level. If the morphological analysis cannot identify the specimen, we will categorize it at the higher taxonomic level.

Zooplankton Collection

We conducted a vertical sampling of zooplankton, commencing at a depth of one meter below the water's surface. The water was filtered using a plankton net with a mesh

Statistical Analysis

Statistical analyzes were performed and visualized in R studio (v. 3.6.2, <http://r-projekt.org>). Spatial and temporal visualization of composition data used bar




Figure 1. The sampling location was Bentar Beach, Probolinggo, East Java. It divided into six station, (PR_1, PR_2, PR_3, PR_4 and PR_5, PR_6).

plots from the ggplot2 package. Data were analyzed by non-parametric statistics and are recommended for this study (Nelson and Eckert, 2007). Kruskal-Wallis test was used to test statistical differences between zooplankton composition between station and month. Kruskal-Wallis test was used to test statistical differences between zooplankton composition, station and month. Besides, Kruskal-Wallis was used to the whale shark sighting between station and month. Pearson correlation tests were carried out to test a relationship between zooplankton composition and individual whale shark encountered (Nelson and Eckert, 2007; Hernández-nava and Álvarez-borrego, 2013; Boldrocchi *et al.*, 2018; De Veaux *et al.*, 2018).

RESULTS AND DISCUSSION

Spatial Variation

These observations' spatial variations refer to the horizontal surface distribution at each station (McManus and Woodson, 2012). We have divided these observations into six observation points, drawing on previous research (Kamal *et al.*, 2016; Himawan, 2017; Syah *et al.*, 2018; Kamal *et al.*, 2020) and information from local residents, particularly whale shark tourism activists in Probolinggo. Based on the Kruskal-Wallis analysis, the appearance of whale sharks at the six stations showed significant differences (chi-square test, $X^2 = 1418.6$, $P < 0.05$). Dunn's follow-up test revealed that almost all stations showed significant differences between one station and another. PR_5 found the most whale sharks at each station, with six individuals, whereas PR_4 and PR_6 did not find any individual whale sharks (Figure 2). We conducted the Pearson correlation test to examine the relationship between the appearance of whale sharks and the spatial abundance of zooplankton. The relationship between whale sharks and zooplankton abundance formed a non-

significant negative correlation ($r = -0.01$, $P > 0.05$).

The existence of whale sharks is influenced by various factors including temperature, the level of disturbance they experience, and food availability (Hacohen-Domene *et al.*, 2006; Rowat and Gore, 2007; Whitehead *et al.*, 2019; Marliana *et al.*, 2018). Tropical coastlines with high temperatures are often an attraction for whale sharks. Various investigations (Compagno, 1984; Yap-dejeto *et al.*, 2013; Ketchum *et al.*, 2013; Whitehead *et al.*, 2019; Maharani *et al.*, 2014; Wulandari *et al.*, 2018) have detected this occurrence, and some have recorded temperatures exceeding 35 °C (Robinson *et al.*, 2013). According to research conducted by Wulandari *et al.* (2018), the temperature range in Probolinggo waters fluctuates between 28.2 to 31.3 °C. The higher water temperature in Probolinggo can be caused by various factors, including the large influx of fresh water from rivers and relatively calm wave conditions (Nuriyanto *et al.*, 2019). The presence of fresh water molecules causes increased friction between salt water and fresh water from rivers, resulting in higher temperatures near the coastline than offshore (Tarigan and Edward, 2000). Apart from that, the remote location of PR_5 station and farthest from tourist activities at Bentar Beach is a contributing factor to the presence of sharks at this station.

The total number of zooplankton identified was twenty-three taxa, consisting of seventeen genera and six identified at the higher taxa level (family, class, or order). Different results from Utojo and Mustofa's (2016) research on zooplankton in Probolinggo waters revealed seven genera. The Kruskal-Wallis test on the composition of zooplankton in Probolinggo shows that there are no significant differences between station (chi-square test, $X^2 = 61.465$, $P > 0.05$). Several studies, such as Mirón *et al.* (2014), Cardenas-Palomo *et al.* (2015), and Witalis *et al.* (2024), support the findings of




Figure 2. Average of zooplankton composition (Bar plot $\pm$ SD) and total of *Rhincodon typus* sighting (black line) recorded for each sampling station in one years.

this study. All these studies demonstrate that there were no significant differences in composition of zooplankton between station in same location. Physical oceanographic factors have no effect on the horizontal spatial distribution compared to the vertical spatial distribution, especially in trophic area (McManus and Woodson, 2012; Azevedo and Bonecker, 2003; Kâ *et al.*, 2012; Mirón *et al.*, 2014). Station PR_4 had the highest average abundance (7.49 ± 2.98 ind/m³), while station PR_3 had the lowest average abundance (1.69 ± 0.31 ind/m³) (Figure 2).

The highest number of taxa identified was at station PR_4, totaling 21 taxa and consisting of sixteen genera and six large groups. Several environmental conditions, such as high levels of nutrients, influence the abundance of zooplankton at station PR_4, which indicated can increase the abundance of phytoplankton as a food for zooplankton (Klais *et al.*, 2016).

The genus *Tintinnopsis* sp. accounted for the highest percentage of zooplankton at six stations, with 33%, followed by the Nauplius group at 31%, and fish eggs at 10%. We most often find the *Tintinnopsis* sp. species at point PR_1 (8.1% of the time), the Nauplius group at point PR_4 (14% of the time), and fish eggs at point PR_4 (Figure 2). Due to its cosmopolitan nature, *Tintinnopsis* sp. frequently inhabits neritic areas that are relatively shallower and undulating (Dolan and Pierce, 2013). This cosmopolitan nature is what causes *Tintinnopsis* sp. and *Stenoseniella* sp. to be the main contributors to spatial abundance formation (Feng *et al.*, 2015). However, this study did not find *Stenoseniella* sp. but found other genera, such as *Favella* sp. and *Leprotintinnus* sp., supporting spatial abundance in Probolinggo waters (Dolan and Pierce, 2013). Environmental variables, particularly nutrition, exhibit a strong correlation with




Figure 3. Percent of zooplankton composition (%) for ten taxon dominant in each station at Bentar Beach, Probolinggo.

tintinnid groups (Feng *et al.*, 2015; Jiang *et al.*, 2016; Xu *et al.*, 2016). Research by Feng *et al.* (2015) and Rakshit *et al.* (2017) suggests that the Tintinnids group, closely related to environmental variables, can serve as a bioindicator to distinguish water quality status. Apart from the tintinnids group, there is a Nauplius group that dominates these waters (Takashi and Uchiyama, 2008). Nauplius is a group that prefers relatively warm temperatures and is rich in nutrients. Nauplius itself consists of several forms, such as copepods, barnacles, crabs, and others. The Nauplius stage has a small and almost identical shape, so this shape will be difficult to identify at the species level (Jungbluth *et al.*, 2017).

Rivers heavily influence Probolinggo Beach, which features relatively calmer waves (Nuriyanto *et al.*, 2019). Mangrove

forests, which encircle these waters, trap organic material input from rivers and the sea itself, thereby increasing the concentration of nutrients in the surrounding waters (Nugroho *et al.*, 2018; Boldrocchi *et al.*, 2020). Anthropogenic activity from rivers greatly influences the beach itself, leading to an increase in the concentration of nutrients entering it. If the numbers are appropriate, coastal areas will become areas with high levels of biodiversity and productive ecosystems (Kemp and Boynton, 2012). While the appearance of whale sharks does not correlate with the abundance of zooplankton in the area, we must consider other factors, like whale shark disturbances, to make the whale shark zone a crucial area for assessing the adequacy of the whale shark population. Establishing whale shark ecotourism areas necessitates the

implementation of management strategies and regulations.

Temporal Variation

The eleven whale sharks were spotted throughout the course of one year. The Kruskal-Wallis analysis revealed that the presence of whale sharks in Probolinggo varied significantly each month (Chi-square test, $X^2 = 81.04$, $p < 0.05$). The Dunn's follow-up test indicates significant variations in the frequency of whale shark sightings throughout the year, with the greatest number observed between the months of March and November. In total, three individual whale sharks were recorded throughout this period (Figure 4). The seasonal occurrence of whale sharks in Probolinggo is consistent with

previous research conducted by Boldrocchi *et al.* (2020), Whitehead *et al.* (2019), Reyes-Mendoza *et al.* (2021), and Cardenas-Palomo *et al.* (2015). The seasonal appearance of it is influenced by various factors, such as the availability of food (Ketchum *et al.*, 2013; Rohner *et al.*, 2015; Marcus *et al.*, 2016). Prior research (Whitehead *et al.*, 2020a; Hernandez-Nava and Alvarez-Borrego, 2013; Boldrocchi *et al.*, 2020; Robinson *et al.*, 2016; Ryan *et al.*, 2017) has investigated the correlation between these variables. Whale sharks in the waters of Probolinggo engage in active surface feeding, which involves swimming on the surface, constantly opening and closing their mouths, and filtering food items through their gill apparatus (Figure 5) (Motta *et al.*, 2010; Cade *et al.*, 2020). The correlation analysis conducted to examine the




Figure 4. Average of zooplankton composition (Bar plot $\pm$ SD) and total of *Rhincodon typus* sighting (black line) recorded for each month in six station, Bentar Beach, Probolinggo.



Figure 5. A whale shark feeding actively at the surface. The whale shark will open the mouth and move actively for filter the water.

relationship between the occurrence of whale sharks and the quantity of zooplankton per month reveals a weak positive connection that is not statistically significant ($r = 0.01$, $P > 0.05$).

The abundance of zooplankton in Probolinggo waters has a temporal pattern. The highest abundance of zooplankton was in March ($8.19 \pm 2.52 \text{ ind/m}^3$) and April ($13.10 \pm 5.89 \text{ ind/m}^3$) (Figure 4). Kruskal-Wallis analysis showed that reporting of zooplankton in each month was significantly different (Chi-square test, $X^2 = 148.61$, $P < 0.05$). DunnTest shows significant differences in December–August, December–June, and December–March. Research by Whitehead *et al.* (2020b) and Reyes-Mendoza *et al.* (2021), which found temporal differences in species composition and reported zooplankton, supports the results of this study. Temperature and water clarity have an influence on the composition of zooplankton each month (Witalis *et al.*, 2024). The same genus and group, namely *Tintinnopsis sp.* (35%), Nauplius (22%), and fish eggs (10%), dominate the percentage of

spatial and temporal distribution of taxa. The most common times to find *Tintinnopsis sp.* are in January (65%), April (45%), and July (25%) (Figure 6).

The Tintinida group has the ability to dominate microzooplankton communities (Fonda Umani and Beran, 2003; Yap-dejeto *et al.*, 2013). Several factors, including temperature, salinity, and water clarity, influence the abundance of the tintinnid group (Wang *et al.*, 2014). Based on research by Rakshit *et al.* (2017), tintinnid community structure has a positive correlation with environmental variables, especially temperature and chlorophyll a. Low levels of water clarity are associated with Chlorophyll A. If water clarity increases, environmental variables such as temperature and concentration of chlorophyll a will be low (Rakshit *et al.*, 2017). The tintinnid group is not the main target of whale sharks in an ecosystem because this group has a small biomass value (Bojanić *et al.*, 2005). However, the tintinnid group is often associated with other microzooplankton groups and has an important role in




Figure 6. Percent of zooplankton composition (%) for each station in Pantai Bentar, Probolinggo.

channeling energy flows in the ecosystem to higher trophic levels (Conway, 2012; Stoecker and Pierson, 2019). Certain months, such as April and July, exhibit the highest reproduction or growth rates, contributing to the abundance of Nauplius and fish eggs in this study (Dar *et al.*, 1970; Turner, 2002). However, we were unable to identify either the Nauplius group or the fish eggs at the genus level, making it impossible to define a dominant level of reproduction or growth. Apart from that, Nauplius crustaceans are abundant because this group has effective movement to avoid predators and the ability to find prey is relatively effective, so the respiration rate in this group is relatively low (Marshall and Orr, 1966; Kiørboe, 2011).

Tintinnopsis sp., Nauplius, and fish eggs dominated the Probolinggo waters in this study both spatially and temporally, but there was no correlation with whale shark appearance. In this study, we conducted a correlation analysis to examine the impact of whale shark appearances on specific genera or groups. The findings showed that among the whole genus and group, *Acartia sp.* has a

positive correlation with the sighting of whale sharks ($r = 0.3$, $P < 0.05$). Other species, such as *Calanus sp.* ($r = 0.1$, $P > 0.05$) and Polychaeta larvae ($r = 0.2$, $P > 0.05$), also had a positive but not significant with the sighting of whale sharks in Probolinggo. Research by Whitehaed *et al.* (2020a) yielded different results, indicating that the copepod group in the waters of Bahia de La Paz, Mexico, did not exhibit a significant correlation, while the "other" group demonstrated a relationship with the appearance of whale sharks. Theoretically, the concentration or density of their food influences the appearance of whale sharks (Heyman *et al.*, 2001; Wilson *et al.*, 2001; de la Parra Venegas *et al.*, 2011; Rowat *et al.*, 2011; Reyes-Mendoza *et al.*, 2021). In areas where whale sharks are most prevalent, their aggregations react to a variety of food sources, including zooplankton blooms (Ketchum *et al.*, 2013; Cárdenas-Palomo *et al.*, 2015; Lavaniegas *et al.*, 2017) and fish spawning (Heyman *et al.*, 2012). Because whale sharks are opportunistic creatures (Pierce and Norman, 2016), they will

congregate in a location that can meet their nutritional needs (Rohner *et al.*, 2015; Boldrocchi *et al.*, 2020). Apart from that, the whale shark's feeding strategy also determines and will be different for each region (Lawson *et al.*, 2019), where in Probolinggo waters the whale shark's feeding type is active surface feeding. Active surface feeding is the way whale sharks eat by opening their mouths while moving on the surface (Motta *et al.*, 2010; Cade *et al.*, 2020). This condition causes the whale shark to move more, potentially increasing the current entering its mouth and influencing the amount of zooplankton it filters. This condition is the same as for baleen whales (Potvin and Werth, 2017; der Hoop *et al.*, 2019).

The abundance of zooplankton and the appearance of whale sharks in Probolinggo are two forms of basic information. The Probolinggo Waters area, renowned for whale shark tourism activities, can utilize this basic information. We can use this data to understand the food availability of whale sharks and to influence their emergence. Therefore, we can create a platform that recommends policies for whale sharks and minimizes their disturbance during foraging events.

CONCLUSION

The temporal variations in the zooplankton composition and whale shark presence are significant. There is no apparent correlation between the zooplankton composition in Probolinggo and whale sharks. Nonetheless, a substantial correlation has been found between the emergence of whale sharks and some taxa, including *Acartia* sp. The zooplankton composition at each of the six observation stations did not significantly differ from one another in the spatial analysis; however, there were distinct variations in the appearance of whale sharks at each station. There is no discernible relationship between the distribution of

zooplankton and the presence of whale sharks.

ACKNOWLEDGMENT

We Are Grateful to team of Whale Shark Indonesia and Marine Biodiversity Laboratory to provide and to help in the research. Thank to National Geographic Society Early Career Grant 2017 to support this research.

REFERENCES

- Bojanić, N., M. Šolić, N. Krstulović, S. Šestanović, I. Marasović, and Ž. Ninčević. 2005. Temporal variability in abundance and biomass of ciliates and copepods in the eutrophicated part of Kaštela Bay (Middle Adriatic Sea). *Helgol. Mar. Res.*, 59(2): 107–120. <https://doi.org/10.1007/s10152-004-0199-x>
- Boldrocchi, G., M. Omar, A. Azzola, and R. Bettinetti. 2020. The ecology of the whale shark in Djibouti. *Aquat. Ecol.*, 54(2): 535–551. <https://doi.org/10.1007/s10452-020-09758-w>
- Boldrocchi, G., Y.M. Omar, D. Rowat, and R. Bettinetti. 2018. First results on zooplankton community composition and contamination by some persistent organic pollutants in the Gulf of Tadjoura (Djibouti) science of the total environment first results on zooplankton community composition and contamination by some p. *Sci. Total Environ.*, 627(February): 812–821. <https://doi.org/10.1016/j.scitotenv.2018.01.286>
- Cade, D.E., J.J. Levenson, R. Cooper, R. de la Parra, D.H. Webb, and A.D. Dove. 2020. Whale sharks increase swimming effort while filter feeding, but appear to maintain high foraging efficiencies. *Journal of Experimental Biology.* 223(11): p.jeb224402.

- <https://doi.org/10.1242/jeb.224402>.
- Cárdenas-Palomo, N., J. Herrera-Silveira, I. Velázquez-Abunader, O. Reyes, and U. Ordoñez. 2015. Distribution and feeding habitat characterization of whale sharks *Rhincodon typus* in a protected area in the north Caribbean Sea. *J. Fish Biol.*, 86(2): 668–686. <https://doi.org/10.1111/jfb.12589>
- Conway, D.V.P. 2012. Marine Zooplankton of Southern Britain Part 1: Radiolaria, Heliozoa, Foraminifera, Ciliophora, Cnidaria, Ctenophora, Platyhelminthes, Nemertea, Rotifera and Mollusca. *Mar. Zooplankt. South. Britain*, (25): 10–139. <http://plymsea.ac.uk/id/eprint/5631>
- Dar, I.A., H.A. Rather, and M.A. Dar. 1970. Dynamics of zooplankton in relation to physico-chemical factors. *Our Nat.*, 7(1): 168–176. <https://doi.org/10.3126/on.v7i1.2567>
- de la Parra Venegas, R., R. Hueter, J.G. Cano, J. Tyminski, J.G. Remolina, M. Maslanka, A. Ormos, L. Weigt, B. Carlson, and A. Dove. 2011. An unprecedented aggregation of whale sharks, *Rhincodon typus*, in Mexican coastal waters of the Caribbean sea. *PLoS One*, 6(4): p.e18994. <https://doi.org/10.1371/journal.pone.0018994>
- De, Veaux, R.D., P.F. Velleman, D.E. Bock, A.M. Vukov, and A.C. Wong. 2005. *Stats: data and models*. Boston: Pearson/Addison Wesley. 652-708 pp.
- Feng, M., W. Zhang, W. Wang, G. Zhang, T. Xiao, and H. Xu. 2015. Can tintinnids be used for discriminating water quality status in marine ecosystems?. *Marine pollution bulletin*. 101(2): 549-555. <https://doi.org/10.1016/j.marpolbul.2015.10.059>.
- Fonda, S. Umani, and A. Beran. 2003. Seasonal variations in the dynamics of microbial plankton communities: First estimates from experiments in the Gulf of Trieste, Northern Adriatic Sea. *Mar. Ecol. Prog. Ser.*, 247: 1–16. <https://doi.org/10.3354/meps247001>
- Graham, R.T., C.M. Roberts, and J.C.R. Smart. 2006. Diving behaviour of whale sharks in relation to a predictable food pulse. *J. R. Soc. Interface*, 3(6): 109–116. <https://doi.org/10.1098/rsif.2005.0082>
- Hacohen-Domene, A., F. Galvan-Magana, and J. Ketchum-Mejia. 2006. Abundance of whale shark (*Rhincodon typus*) preferred prey species in the southern Gulf of California, Mexico. *Cybiurn*, 30(4): 99–102. <http://pascal-francis.inist.fr/vibad/index.php?action=getRecordDetail&idt=18651430>
- Hernandez-Nava, M.F. and S. Alvarez-Borrego. 2013. Zooplankton in a whale shark (*Rhincodon typus*) feeding area of Bahia de los Angeles (Gulf of California). *Hidrobiologica*, 23(2): 198–208. <https://hidrobiologica.izt.uam.mx/hidrobiologica/index.php/revHidro/article/view/638>
- Heyman, W.D., R.T. Graham, B. Kjerfve, and R.E. Johannes. 2001. Whale sharks *Rhincodon typus* aggregate to feed on fish spawn in Belize. *Mar. Ecol. Prog. Ser.*, 215: 275–282. <https://doi.org/10.3354/meps215275>
- Himawan, M.R. 2017. Komposisi hiu paus berdasarkan jenis kelamin, ukuran, perilaku kemunculan serta ancaman keberadaannya di Perairan Indonesia [tesis]. Bogor : Institut Pertanian Bogor. Page: 11-12
- Himawan, M.R., C. Tania, B.A.Noor, A. Wijonarno, B. Subhan, and H. Madduppa. 2015. Sex and size range composition of whale shark (*Rhincodon typus*) and their sighting behaviour in relation with fishermen lift-net within Cenderawasih Bay National Park, Indonesia. *Aquaculture, Aquarium, Conservation and Legislation*, 8(2): 123-133.

- <http://www.bioflux.com.ro/aac1>
- Hoffmayer, E.R., J.S. Franks, W.B. Driggers, K.J. Oswald, and J.M. Quattro. 2007. Observations of a feeding aggregation of whale sharks, *Rhincodon typus*, in the North Central Gulf of Mexico. *Gulf Caribb. Res.*, 19(2): 69–73. <https://doi.org/10.18785/gcr.1902.08>
- Jiang, Y., G. Xu, and H. Xu. 2016. Use of multivariate dispersion to assess water quality based on species composition data. *Environmental Science and Pollution Research*. 23: 3267-3272. <https://doi.org/10.1007/s11356-015-5583-3>
- Jungbluth, M.J., K.E. Selph, P.H. Lenz, and E. Goetze. 2017. Species-specific grazing and significant trophic impacts by two species of copepod nauplii, *Parvocalanus crassirostris* and *Bestiolina similis*. *Marine Ecology Progress Series*. 572: 57-76. <https://doi.org/10.3354/meps12139>
- Kamal, M.M., D. Ardania, and M.T. Hartanto. 2020. Abundance and composition of potential food items of whale shark (*Rhincodon typus* Smith, 1828) in Probolinggo waters, East Java Province. *Eco. Env. Cons.*, 26(2007): 37–41. <http://www.envirobiotechjournals.com/EEC/v26JuneSuppl2020/EEC-7.pdf>
- Kamal, M.M., Y. Wardiatno, and N.S. Noviyanti. 2016. Habitat conditions and potential food items during the appearance of whale sharks (*Rhincodon typus*) in Probolinggo waters, Madura. *The 4th International Whale Shark Conference*, Hamad bin Khalifa University Press, 2016(2): 27p. <https://doi.org/10.5339/qproc.2016.iws.c4.27>
- Ketchum, J. T., F. Galván-Magaña, and AP. Klimley. 2013. Segregation and foraging ecology of whale sharks, *Rhincodon typus*, in the southwestern Gulf of California. *Environmental Fish Biology*, 96: 779–755. <https://doi.org/10.1007/s10641-012-0071-9>
- Kjørboe, T. 2011. What makes pelagic copepods so successful? *J. Plankton Res.*, 33(5): 677–685. <https://doi.org/10.1093/plankt/fbq159>
- Klais, R., M. Lehtiniemi, G. Rubene, A. Semenova, P. Margonski, A. Ikauniece, M. Simm, A. Põllumäe, E. Grinienė, K. Mäkinen, et al.,., 2016. Spatial and temporal variability of zooplankton in a temperate semi-enclosed sea: implications for monitoring design and long-term studies. *J. Plankton Res.*, 38(3): 652–661. <https://doi.org/10.1093/plankt/fbw022>
- Lavaniegos, B., G. Heckel, and P. de Guevara. 2017. Seasonal variability of copepods and cladocerans in Bahía de los Ángeles (Gulf of California) and importance of *Acartia clausi* as food for whale sharks. *Ciencias Mar.*, 38(2012): 11–30. <https://doi.org/10.7773/cm.v38i1A.2017>
- Lawson, C.L., L.G. Halsey. G.C. Hays, C.L. Dudgeon, N.L. Payne, M.B. Bennett, C.R. White, and A.J. Richardson. 2019. Powering Ocean Giants : The energetics of Shark and Ray megafauna. *Trends Ecol. Evol.* 34: 1009–1021. <http://dx.doi.org/10.1016/j.tree.2019.07.001>
- Marliana, S.N., M. Bataona, and E.N. Ihsan. 2018. Zooplankton communities in Cenderawasih Bay National Park, West Papua: can their composition be used to predict whale shark *Rhincodon typus* Smith, 1828 appearance frequencies?. *IOP Conference Series: Earth and Environmental Science*, 139(1): 012012. <https://doi.org/10.1088/1755-1315/139/1/012012>
- Marshall, S.M. and A.P. Orr. 1966. Respiration and feeding in some small copepods. *J. Mar. Biol. Assoc. United Kingdom.*, 46(3): 513–530.

- <https://doi.org/10.1017/S0025315400033312>
- McKinney, J.A., E.R. Hoffmayer, W. Wu, R. Fulford, and J.M. Hendon. 2012. Feeding habitat of the whale shark *Rhincodon typus* in the northern Gulf of Mexico determined using species distribution modelling. *Mar. Ecol. Prog. Ser.*, 458: 199–211. <https://doi.org/10.3354/meps09777>
- Meekan, M.G., S.N. Jarman, C. McLean, and M.B. Schultz. 2009. DNA evidence of whale sharks (*Rhincodon typus*) feeding on red crab (*Gecarcoidea natalis*) larvae at Christmas Island, Australia. *Mar. Freshw. Res.*, 60(6): 607–609. <https://doi.org/10.1071/MF08254>
- Nelson, J.D. and S.A. Eckert. 2007a. Foraging ecology of whale sharks (*Rhincodon typus*) within Bahía de Los Angeles, Baja California Norte, México. *Fish. Res.*, 84(1): 47–64. <https://doi.org/10.1016/j.fishres.2006.11.013>
- Nour, and N.M. El-Din. 2001. Ecological investigation of the tintinnid community along the coastal waters of Alexandria, Egypt. *Egyptian Journal of Aquatic Biology and Fisheries*, 5(3): 165–177. https://ejabf.journals.ekb.eg/article_1698.html
- https://ejabf.journals.ekb.eg/article_1698_dcf6ad7ef870e2e5ec148a8e0f098c06.pdf
- Nugroho, T.T., M. Najib, and K. Kirbrandoko, K. 2018. Determination of competitiveness based on core competency analysis (case study on mangrove ecotourism in East Java). *Matrik J. Manajemen, Strateg. Bisnis Dan Kewirausahaan* 12(1): 25–32. <https://doi.org/10.24843/MATR-12-1.01.P03>
- Nuriyanto, M.Z., F.A. Firmansyah, and I. Prasetyo. 2019. Analisis perubahan bentang geomorfologi pantai Bentar Kabupaten Probolinggo. *Maj. Pembelajaran Geogr.*, 2(1): 99–109. <<https://jurnal.unej.ac.id/index.php/PGEO/article/view/11523>>
- Oxford University. 2017. Marine plankton: A practical guide to ecology, methodology, and taxonomy. Castellani C, and Edwards M. eds. Oxford University Press. 181–609 pp.
- Pierce, R.W. and J.T. Turner. 1993. Global biogeography of marine tintinnids. *Mar. Ecol. Prog. Ser.*, 94(1): 11–26. <https://doi.org/10.3354/meps094011>.
- Pierce, S.J., B. Norman. 2016. *Rhincodon typus*. In: The IUCN Red List of Threatened Species 2016. e.T19488A2365291. <https://doi.org/10.2305/IUCN.UK.2016-1.RLTS.T19488A2365291.en>.
- Potvin, J., A.J. Werth. 2017. Oral cavity hydrodynamics and drag production in Balaenid whale suspension feeding. *PLoS ONE* 12: e0175220. <https://doi.org/10.1371/journal.pone.0175220>.
- Rakshit, D., K. Murugan, J.K. Biswas, K.K. Satpathy, P.S. Ganesh, and S.K. Sarkar, S.K. 2017. Environmental impact on diversity and distribution of tintinnid (Ciliata: Protozoa) along Hooghly Estuary, India: a multivariate approach. *Regional Studies in Marine Science*. 12: 1–10. <https://doi.org/10.1016/j.rsma.2017.02.007>.
- Reyes-Mendoza, O., N. Cárdenas-Palomo, J. Herrera-Silveira, E. Mimila-Herrera, J. Trujillo-Córdova, X. Chiappa-Carrara, and D. Arceo-Carranza. 2021. Quantity and quality of prey available for the whale shark (*Rhincodon typus*, Smith 1828) at the Mexican Caribbean aggregation site. *Regional Studies in Marine Science*, 43: 101696. <https://doi.org/10.1016/j.rsma.2021.101696>.
- Robinson, D.P., M.Y. Jaidah, S. Bach, K. Lee, W. Jabado, C.A. Rohner, A. March, and S. Caprodossi. 2016.

- Population structure, Abundance and movement of Whale Sharks in the Arabian Gulf and the Gulf of Oman. *PLoS ONE*. 11(6), <http://dx.doi.org/10.1371/journal.pone.0158593>
- Rohner, C.A., A.J. Armstrong, S.J. Pierce, C.E.M. Prebble, E.F. Cagua, J.E.M. Cochran, M.L. Berumen, and A.J. Richardson. 2015. Whale sharks target dense prey patches of sergestid shrimp off Tanzania. *J. Plankton Res.* 37: 352–362. <http://dx.doi.org/10.1093/plankt/fbv010>
- Rowat, D., K. Brooks, A. March, C. McCarten, D. Jouannet, L. Riley, G. Jeffreys, M. Perri, M. Vely, and B. Pardigon. 2011. Long-term membership of whale sharks (*Rhincodon typus*) in coastal aggregations in Seychelles and Djibouti. *Mar. Freshw. Res.*, 62(6): 621–627. <https://doi.org/10.1071/MF10135>
- Rowat, D. and U. Engelhardt. 2007. Seychelles: A case study of community involvement in the development of whale shark ecotourism and its socio-economic impact. *Fish. Res.*, 84(1): 109–113. <https://doi.org/10.1016/j.fishres.2006.11.018>
- Rowat, D. and M. Gore. 2007. Regional scale horizontal and local scale vertical movements of whale sharks in the Indian Ocean off Seychelles. *Fish. Res.*, 84(1): 32–40. <https://doi.org/10.1016/j.fishres.2006.11.009>
- Ryan, J.P., J.R. Green, E. Espinoza, and A.R. Hearn. 2017. Association of whale sharks (*Rhincodon typus*) with thermobiological frontal systems of the eastern tropical Pacific. *PLoS ONE*. 12(8). <http://dx.doi.org/10.1371/journal.pone.0182599>
- Stoecker, D. and J. Pierson. 2019. Predation on protozoa: Its importance to zooplankton revisited. *J. Plankton Res.*, 41(4): 367–373. <https://doi.org/10.1093/plankt/fbz027>
- Syah, A.F., M. Musrifah, and H. Cahyono. 2018. Pemodelan daerah potensial kemunculan hiu paus (*Rhincodon typus*) menggunakan data penginderaan jauh di perairan probolinggo, jawa timur. *J. Penelit. Perikan. Indones.*, 24(3): 209. <https://doi.org/10.15578/jppi.24.3.2018.209-216>
- Turner, J.T. 2002. Zooplankton fecal pellets, marine snow and sinking phytoplankton blooms. *Aquat. Microb. Ecol.*, 27(1): 57–102. <https://doi.org/10.3354/ame027057>
- Utojo, M.A. 2016. Struktur Komunitas Plankton pada Tambak Intensif dan Tradisional Kabupaten Probolinggo, Provinsi Jawa Timur. *Jurnal Ilmu dan Teknologi Kelautan Tropis*, 8(1): 269.
- van der Hoop, J. M., A.E. Nousek-McGregor, D.P. Nowacek, S.E. Parks, P. Tyack, and P.T. Madsen. 2019. Foraging rates of ram-filtering North Atlantic right whales. *Funct. Ecol.* 33: 1290-1306. <https://doi.org/10.1111/1365-2435.13357>
- Wang, Y., W. Zhang, Y. Lin, L. Zheng, W. Cao, and J. Yang. 2014. Spatial and seasonal variations of large tintinnid ciliates in Shenhui Bay of China. *Oceanol. Hydrobiol. Stud.*, 43(3): 292–302. <https://doi.org/10.2478/s13545-014-0144-9>
- Wilson, S.G. 2002. A whale shark feeding in association with a school of giant herring at Ningaloo Reef, Western Australia. *J. R. Soc. West. Aust.*, 85(1): 43–44. <https://www.proquest.com/scholarly-journals/whale-shark-feeding-association-with-school-giant/docview/762205106/se-2?accountid=201395>

- Wilson, S.G., J.G. Taylor, and A.F. Pearce. 2001. The seasonal aggregation of whale sharks at Ningaloo Reef, Western Australia: Currents, migrations and the El Niño/ Southern Oscillation. *Environ. Biol. Fishes.*, 61(1): 1–11. <https://doi.org/10.1023/A:1011069914753>
- Witalis, B., A. Iglukowska, M. Ronowicz, A. Weydmann-Zwolicka, and P. Kukliński. 2024. The zooplankton community of Baltic Sea ports: diversity and seasonal dynamics. *Oceanologia*. <https://doi.org/10.1016/j.oceano.2024.02.001>.
- Whitehead, D.A., D. Petatán Ramirez, D. Olivier, R. González-Armas, F. Pancaldi, and F. Galvan-Magana. 2019. Seasonal trends in whale shark *Rhincodon typus* sightings in an established tourism site in the Gulf of California, Mexico. *Journal of Fish Biology*, 95(3): 982–984. <https://doi.org/10.1111/jfb.14106>
- Whitehead, D.A., U. Jakes-Cota, F. Pancaldi, F. Galván-Magaña, and R. González-Armas. 2020a. The influence of zooplankton communities on the feeding behavior of whale shark in Bahia de La Paz, Gulf of California. *Revista mexicana de biodiversidad*, 91: e913054. <https://doi.org/10.22201/ib.20078706e.2020.91.3054>.
- Whitehead, D.A., U. Jakes-Cota, F. Galván-Magaña, F. Pancaldi, and R. González-Armas. 2020b. Composition and abundance of macro zooplankton in the coastal waters off the El Mogote Sandbar. *Hidrobiológica*. 30(1): 21-27. <https://doi.org/10.24275/uam/izt/dcbs/hidro/2020v30n1/whitehead>.
- Xu, H., Y. Jiang, and G. Xu. 2016. Identifying functional species pool of planktonic protozoa for discriminating water quality status in marine ecosystems. *Ecological indicators*. 62: 306-311. <https://doi.org/10.1016/j.ecolind.2015.10.068>
- Yap-dejeto, L., A. Cera, J. Labaja, J.D. Palermo, and A. Ponzo. 2013. Observations of Microzooplankton in the Vicinity of Whale Shark *Rhincodon typus* Aggregation Sites in Oslob , Cebu and Pintuyan , S . Leyte , Philippines. (December):61–77. <https://www.lamave.org/s/Yap-dejetoetal2018ZoopandwhalesharksPJNS.pdf>

