

Genetic Diversity of the Structure of HSP70 Gene in Kampung Unggul Balitbangtan (KUB), Walik, and Kate Walik Chickens

A. Aryani^a, D. D. Solihin^{b,*}, C. Sumantri^c, R. Afnan^c, & T. Sartika^d

^aDepartment of Biology Education, Faculty of Mathematics and Natural Science Education,
Indonesia University of Education,
Jalan Dr. Setiabudi No. 229 Bandung 40154, Indonesia

^bDepartment of Biology, Faculty of Mathematics and Natural Sciences, IPB University (Bogor Agricultural University)

^cDepartment of Animal Production and Technology, Faculty of Animal Science,
IPB University (Bogor Agricultural University)

Jalan Agatis, Kampus IPB Darmaga Bogor 16680, Indonesia

^dIndonesian Research Institute for Animal Production

Jalan Veteran III, PO. BOX 221 Ciawi Bogor 16002, Indonesia

*Corresponding author: dduryadi@yahoo.com

(Received 15-05-2018; Revised 05-04-2019; Accepted 25-04-2019)

ABSTRACT

Our research was designed to identify the sequence variations of heat shock protein 70 (HSP70) gene in three breeds of native Indonesian chickens, Kampung Unggul Balitbangtan (KUB) chicken, Walik chicken, and Kate Walik chicken. Total DNA was isolated from the whole blood using a DNeasy blood and tissue kit. The HSP70 gene was amplified and sequenced from 94 chickens using PCR. The amplification product was 787 bp long, consisting of a 210 bp promoter region, a 112 bp long 5'untranslated region (UTR), and a 465 bp protein coding region. Our KUB, Walik, and Kate Walik chicken HSP70 gene sequence alignments express genetic diversity in the promoter region (insertions and deletions), 5'UTR (deletions and nucleotide substitutions), and at the beginning of the coding region (nucleotide substitutions). Four haplotypes, H1, H2, H3, and H4, were identified in the HSP70 gene protein coding region. The haplotype H2 was found in all three chickens, while H4 was only found in Walik chicken. The H4 is a novel haplotype which never reported before. Based on a median-joining network analysis, H4 is a haplotype produced by mutations at two specific sites (g.370A>G and g.388C>G) in the protein coding region of the HSP70 gene of the chicken. It could be concluded that Walik chicken can be used as a standard for heat stress genotyping in Indonesian local chickens, because it has complete HSP70 gene haplotypes.

Keywords: HSP70 gene; Indonesian native chicken; polymorphism; haplotype; PCR-Sequencing

INTRODUCTION

Indonesia has a high diversity of local chicken breeds, which have the potential to be developed as a national food security source, but their utilizations are not optimal. So far, 31 breeds of local chickens have been identified (Nataamijaya, 2010). Indonesian local chickens, aside from being kept as broilers and egg-laying hens, are also kept as domestic companions, serving as yard decorators, the source of neighborly complaints, and the provider of ritual needs, or merely to provide gratification through melodious voices (Sartika *et al.*, 2016). One of Indonesia's local chickens is the Walik chicken having curly body feathers (frizzle). Adult Walik chickens are quite resistant to weather changes. However, Walik chickens have a rather limited population, existing only in certain regions, and are, therefore, rare local chickens that need to be further explored (Sartika *et al.*, 2016). In addition to local chickens, Kampung Unggul Balitbangtan (KUB) chickens, which are the result of six generations of selection using female

chickens from Cianjur, Depok, Majalengka, and Bogor regions, have several advantages. These include the high egg production and the majority do not brood (Minister of Agriculture Decree No. 274/Kpts/SR.120/2/2014).

Poultry does not have sweat glands, and almost all of the poultry body is covered by feathers. Therefore, dissipation of body heat to the environment is difficult, and poultry is quite susceptible to heat stress (Tamzil, 2014). High environmental temperature has negative impacts on physiological conditions (metabolic and hormonal activity, and body temperature control) and chicken productivity. Heat stress is a primary poultry problem worldwide, especially in broiler and egg-laying chickens. Heat stress can occur anytime when ambient temperature exceeds the limit of comfort zone (>28 °C). Heat stress in chickens can cause growth disruption, decreased egg quality, and death.

Heat shock protein 70 (HSP70), among others, is produced by organisms to overcome heat stress. The HSP70 gene works as a chaperone, the duty of which is to regulate the refolding of proteins properly, to

help protect cells from damage caused by heat stress (Tkáčová & Angelovičová, 2012). The HSP70 gene in chicken, *Gallus gallus*, is located in the fifth chromosome and consists of only one exon with a coding-region length of 1,905 bp (Morimoto *et al.*, 1986). The complete sequence of the chicken HSP70 gene is available in GenBank (J02579). Research on the polymorphisms of HSP70 gene between breeds of chickens has progressed based on the sequence of HSP70 gene of chickens. One such study identified polymorphisms in the coding region of the HSP70 genes in broiler chickens of different breeds at positions g.370A>G (identical to g.258A>G from the start codon) and g.388C>G (identical to g.276C>G from the start codon). Three of the detected alleles (hsp70-1, hsp70-2, and hsp70-3) were associated with the nature of resistance to heat stress, and therefore, it's potential to be used as a candidate gene for heat stress (Mazzi *et al.*, 2003). The results of this study have encouraged additional research in the other chickens: Indonesian local chicken (Tamzil *et al.*, 2013a), Taiwanese local chicken (Liang *et al.*, 2016), and Thailand local chicken (Duangjinda *et al.*, 2017).

The purpose of this study was to obtain data on the diversity of HSP70 genes in Indonesian local chickens: KUB, Walik, and Kate Walik chickens. We identified the genetic variability of the chicken HSP70 gene promoter, 5'UTR, and coding region. These data are essential, both as an informational database, especially in response to a high environmental temperature during summer, and as a basis for the development of heat-resistant chicken strains. In this study, we used two different feather structures. KUB chicken has a normal feather, but Walik chicken and Kate Walik chicken have frizzle feathers. The frizzle (F) is an autosomal and incompletely dominant gene, which has a nonframeshift deletion in KRT75 (Ng *et al.* 2012). The F gene may reduce the heat insulation of feathers by curling and reducing their sizes. Many reports have proved that frizzle (F) and naked neck (Na) are well known as major marker genes affecting heat stress tolerance, resistant to diseases, and better

immune system (Fathi *et al.* 2013, 2014). Several tropical countries have introduced these genes in breeding programs to improve performance and adaptability of chickens to hot ambient temperatures (Fathi *et al.* 2013, 2014; Carabaño *et al.* 2019). Recently, supplementation of vitamin and mineral are used as antioxidants to reduce the negative effects of heat stress (Rayani *et al.* 2017; Mohamed *et al.* 2019).

MATERIALS AND METHODS

Sample Collection and DNA Isolation

In this study, we used 94 chickens from three breeds of native Indonesian chickens: KUB, Walik, and Kate Walik chickens (Table 1). Specifically, the KUB chicken sampling fulfilled the requirements of the Balitbangtan ethics code, registration number Balitbangtan/Balitnak/A/01/2018 in 2018. Chicken DNA material comes from the whole blood taken from the wing vein (\pm 0.5–1.0 mL). Genome DNA extraction and purification were carried out using a DNeasy blood and tissue kit based on the spin column procedure (Qiagen, Venlo, Netherlands).

Amplification and Sequencing

DNA amplification was performed using the polymerase chain reaction (PCR) with a primary pair of forward, 5'-CGATCTGGCTGCAATCTACG-3' (Gan *et al.*, 2015) and reverse, 5'-CTGGGAGTCGTTGAAGT AAGCG-3' (Mazzi *et al.*, 2003) primers based on the chicken HSP70 gene (Morimoto *et al.*, 1986). The amplified PCR target product is 787 bp long, consisting of a 210 bp promoter region, a 112 bp 5' untranslated region (UTR), and a 465 bp coding region (Figure 1). The PCR mixture consisted of 12.5 μ L of GoTaq Green Master Mix (Promega, Madison, Wisconsin) containing 2x GoTaq Green reaction buffer, 400 μ M of each of four deoxynucleoside triphosphates, and

Table 1. Number of chicken samples analyzed

Breed	n	Male	Female	Management system	Source
Kampung Unggul Balitbangtan (KUB)	50	20	30	Intensive (battery cage)	Indonesian Research Institute for Animal Production, Bogor
Walik	41	4	37	Semi Intensive	Private collection
Kate Walik	3	1	2	Semi Intensive	Private collection




Figure 1. Amplification target of the HSP70 gene in chicken (J02579) as a reference. The arrow shows the attachment of the primer: the forward primer attaches at position -210 (promoter region), and the reverse primer attaches at position +577 (protein coding region) of the HSP70 gene of chicken.

3 mM MgCl₂, 1 µL of each primer HSP70 forward and HSP70 reverse (10 µM), 1 uL DNA template (100 ng), and nuclease-free water to final volume of 25 µL. The PCR program consisted of an initial denaturing at 95°C for 5 minutes, followed by 35 cycles of denaturation at 95°C for 30 seconds, annealing at 58°C for 45 seconds, and elongation at 72°C for 1 minute. The amplification process was terminated by extension at 72°C for 5 minutes. PCR products were analyzed by electrophoresis on an agarose gel, stained with ethidium bromide, and then sequenced with ABI® PRISM bigdye terminator cycle sequencing kit v3.1 by First BASE Laboratories (Selangor, Malaysia).

Data Analysis

Chromatogram nucleotide sequencing results, both forward and reverse, were edited using Bioedit (Hall, 2011). The edited nucleotides were then aligned with the reference chicken HSP70 gene from National Center for Biotechnology Information (NCBI) GenBank using Clustal W implemented in Molecular Evolutionary Genetic Analysis (MEGA) version 6 software (Tamura *et al.*, 2013). Similarity searching with our HSP70 gene coding sequences was performed with the Basic Local Alignment Search Tool-nucleotide application (BLASTn) at NCBI. Reconstruction of haplotype networks was made to explore further evolutionary relationships between nucleotide substitutions using Population Analysis with Reticulate Trees (PopART) software (Leigh & Bryant, 2015). Analysis of potential CpG island in the chicken HSP70 gene was detected by MethPrimer (<http://www.urogene.org/methprimer>).

RESULTS

Amplification of the Chicken HSP70 Gene

Partial amplification of HSP70 gene in KUB, Walik, and Kate Walik chickens produced 787 bp DNA fragments, as expected (Figure 2). The sequenced DNA fragments, after being aligned with the reference chicken HSP70 gene, were found to consist of a 210 bp promoter region, a 112 bp 5'UTR, and a 465 bp coding region.

Analysis of the Chicken HSP70 Gene

Promoter Region. The results of sequencing analysis showed that nucleotide variations were found at posi-

tions -103, -99, -11, -6, and -5 in the promoter region of the KUB, Walik, and Kate Walik chicken HSP70 genes compared to reference chicken (Table 2). However, the KUB, Walik, and Kate Walik chickens had identical nucleotide sequences of promoter region of HSP70 gene. Two deletions and three insertions were found in the KUB, Walik, and Kate Walik chicken HSP70 gene partial promoter region sequence alignment, regarding to chicken HSP70 gene (Figure 3). The deletions indicated the loss of a guanine (G) base at position -103, and the loss of a cytosine (C) base at location -99. In addition to these deletions, there were the insertions of two G bases at positions -11 and -6, and the inclusion of one C base at position -5.

The promoter region of chicken HSP70 gene showed consensus-motif sequences consisted of a TATA box, Heat Shock Elements 1 and 2 (HSE 1 and HSE 2), Specificity Protein 1 (SP 1), and CAAT box (Figure 3). The primary recognition element in the promoter region of the chicken HSP70 gene was a TATA box consisted of the sequence 5'-TATAAA-3', found at position -24 through -19. Another recognition element in the promoter region of chicken HSP70 gene was HSE, consisted of HSE 1 with a sequence of 5'-CTGGCAGGTTCCAG-3', and the HSE 2 with a sequence of 5'-CCTTAGCGTTCTGGC-3'. The two heat shock elements overlapped at positions -55 through -42 in HSE 1 and at locations -65 through -51 in HSE 2. The other recognition elements found in the promoter region of chicken HSP70 gene was SP 1 (a type of GC box), with the sequence of 5'-GGGCGG-3', located at position -135 through -130. There were two CAAT box elements with complementary nucleotide sequences of 5'-ATTG-3' found at location -152 through -149, and -71 through -68.

5'UTR. Our sequencing and alignment of the 5'UTR of HSP70 gene in KUB, Walik, and Kate Walik chickens

Table 2. Nucleotide variation in the promoter region of the HSP70 gene in Kampung Unggul Balitbangtan (KUB), Walik, and Kate Walik chickens with respect to the chicken reference sequence

Breed code	n	Nucleotide position				
		-103	-99	-11	-6	-5
Ref. chicken	1	G	C	-	-	-
KUB	50	-	-	G	G	C
Walik	41	-	-	G	G	C
Kate Walik	3	-	-	G	G	C




Figure 2. Visualization of the HSP70 gene PCR products in KUB chicken (a) and Walik chicken (b). M= 1 Kb of DNA Ladder, 1-6= PCR product.

with reference to chicken HSP70 gene sequence showed two deletions and one transition (Figure 4). Two A bases were deleted, one at position +32 and the other at +97. Aside from these deletions, one transition A to G was found at location +44. These nucleotide variations, found at positions +32, +44, and +97 in the 5'UTR, produced two haplotypes (Table 3). The first haplotype was found in KUB (n= 48), Walik (n= 38), and Kate Walik chickens (n= 3), while the second haplotype was only found in KUB (n= 2) and Walik chickens (n= 3). One of these specific nucleotide substitutions in the 5'UTR (g.44A>G) was only found in two KUB chickens and three Walik chickens.

Protein coding region. Four mutations could be seen in our coding region alignment (465 bp) at positions +370, +388, +533, and +537. The four mutations consisted of one transition and three transversions (Table 4). The transition occurred at position g.370A>G (identical to

position g.258A>G from the start codon). The three transversions occurred at positions g.388C>G (identical to position g.276C>G from the start codon), g.533G>C (identical to position g.421G>C from the start codon), and g.537C>A (identical to position g.425C>A from the start codon).

Based on the two mutations (g.370A>G and g.388C>G) found in the HSP70 coding region of KUB, Walik, and Kate Walik chickens, four haplotypes were proposed (Figure 5). We named the four haplotypes Haplotype 1 (H1), Haplotype 2 (H2), Haplotype 3 (H3), and Haplotype 4 (H4). All four haplotypes were occurred in Walik chickens. KUB chickens had three of the haplotypes (H1, H2, and H3), while Kate Walik chickens only had one haplotype (H2). H2 was the most common haplotype and was found in KUB (n= 27), Walik (n= 22), and Kate Walik chickens (n= 3). H1 was found in KUB (n= 7) and Walik chickens (n= 12), while H3 was also found in KUB (n= 16) and Walik chickens (n= 3).



Figure 3. Promoter region sequences of the HSP70 gene in Kampung Unggul Balitbangtan (KUB), Walik, and Kate Walik chickens aligned to the chicken reference sequence. Underlined: forward primer site; bold: consensus motif sequences of CAAT, SP 1, HSEs, and a TATA box (Morimoto *et al.*, 1986); hyphens: nucleotide base deletions.




Figure 4. Alignment of two haplotypes in 5'UTR of the HSP70 gene in Kampung Unggul Balitbangtan (KUB), Walik, and Kate Walik chickens with the chicken reference sequence. Hyphens: nucleotide base deletions; in box: transversion.

Conversely, H4 was only found in Walik chickens (n= 4).

We found that the wild haplotype of HSP70 gene in KUB and Walik chickens was H1, based on our HSP70 haplotype reconstruction using an Median-Joining network (Figure 6). The mutation at position g.388C>G produced H2, while the mutation at position g.370A>G produced H3. The new haplotype, H4, was only found in Walik chickens (n= 4), produced by the mutations at two specific coding regions at positions g.370A>G and g.388C>G of the chicken HSP70 gene.

CpG island. The existence of CpG island could be identified by using MethPrimer. The CpG island was mostly found in the promoter and the first parts of exon in several genes, especially in the housekeeping gene. Only one CpG island with 33 CpG sites was found in the HSP70 gene promoter and exon of KUB, Walik, and Kate Walik chickens (Figure 7). CpG island had 376 bp long, and 61.7% of G+C (Table 5).

DISCUSSION

The promoter region of chicken HSP70 gene is upstream (5'-flanking) of the protein coding sequence and controls the regulation of the transcription process through the binding to various transcription factors that recognize specific DNA motif, i.e. TATA box, HSE, and SP 1 (Morimoto *et al.*, 1986; Mazzi *et al.*, 2003). The TATA box (Goldberg-Hogness box) is generally located around 30 nucleotides upstream of the initiation of transcription within the promoter region in eukaryotic cells, and plays a role in directing RNA polymerase II during transcription. HSE is a specific DNA sequence responsible for the activation of HSP70 gene transcription (Pelham, 1982). HSE elements are binding sites for heat shock [transcription] factors (HSFs) (Pelham, 1982).

The HSP70 gene is considered as an ideal biological marker for heat stress in livestock (Archana *et al.*, 2017). The induction of HSP70 gene expression occurs when stress induces and activates transcription factors, such HSF, to bind to HSE. The binding of HSF to HSE allows the transcription of the HSP70 gene by RNA polymerase II (Akerfelt *et al.*, 2010). Vertebrates have four transcription factors for heat shock gene i.e., HSF1, HSF2, HSF3, and HSF4. Aves have three HSFs, with HSF1 and HSF2 being homologous to transcription factors counterparts

in mammals, whereas HSF3 is a specific transcription factor to aves (Fujimoto & Nakai, 2010). The expression of chicken HSF3 transcription factor is nearly stable throughout the development of various tissues in chicken. HSF2 and HSF4 are present in normal, unstressed cells, playing key roles in response to various biological processes, including immune activation and cell differentiation (Akerfelt *et al.*, 2010). Therefore, HSF1 and HSF3 are likely involved in heat stress response.

In the present study, the analysis of the HSP70 gene in KUB, Walik, and Kate Walik chickens showed the heterogeneity in the promoter region. Nucleotide variations found in the promoter region of the HSP70 gene are caused by the insertions and deletions of nucleotides at positions -103, -99, -11, -6, and -5 that are also found in broiler and naked neck chickens (Mazzi *et al.*, 2003).

The analysis of the 5'UTR of HSP70 gene in KUB, Walik, and Kate Walik chickens detected nucleotide variations occurred at positions +32, +44, and +97. One specific nucleotide substitution in the 5'UTR (g.44A>G) is only found in two KUB chickens and three Walik Chickens (Table 3). When combined with the specific nucleotide substitutions in the protein coding region, these nucleotide substitutions are included in our Haplotype 1 (H1).

Research on the genetic effect of nucleotide substitution in g.69A>G, SNP (single nucleotide polymorphism) in the 5'-flanking region and its association with thermotolerance trait show that only one genotype (GG) proved useful as a marker of heat stress resistance in local Chinese chickens (Gan *et al.*, 2015; Chen *et al.*, 2016). The nucleotide g.69A>G transition reported by Gan *et al.* (2015) and Chen *et al.* (2016) is identical to the position of g.44A>G transition in the 5' UTR of the HSP70 gene of chicken (Morimoto *et al.*, 1986). It is because the number reported by Gan *et al.* (2015) and Chen *et al.* (2016) is based on translational start and the number reported by Morimoto *et al.* (1986) is based on transcriptional start.

A 5'UTR is upstream of the protein-coding sequence in many proteins and is important in regulating gene expression (Araujo *et al.*, 2012; Leppek *et al.* 2018). Several polymorphisms have been reported in the 5'UTR of the bovine HSP70 (bHSP₇₀) gene in Pasundan cattle. Only two of these SNPs, g.1117G>A and g.1125A>C,

Table 3. Haplotypes and nucleotide site variation in the 5'UTR of HSP70 gene in Kampung Unggul Balitbangtan (KUB), Walik, and Kate Walik chickens aligned to the chicken reference sequence

Breed	n	Nucleotide position			Haplotype
		+32	+44	+97	
Ref. chicken	1	A	A	A	-
KUB	48	-	A	-	1
	2	-	G	-	2
Walik	38	-	A	-	1
	3	-	G	-	2
Kate Walik	3	-	A	-	1

Table 4. Haplotypes and mutations in the HSP70 gene protein coding region of Kampung Unggul Balitbangtan (KUB), Walik, and Kate Walik chickens compared with the chicken reference sequence

Breed	N	Nucleotide position				Haplotype
		+370	+388	+533	+537	
Ref. chicken	1	A	C	G	C	-
KUB	7	A	C	C	A	1
	27	A	G	C	A	2
	16	G	C	C	A	3
	12	A	C	C	A	1
Walik	22	A	G	C	A	2
	3	G	C	C	A	3
	4	G	G	C	A	4
	4	G	G	C	A	4
Kate Walik	3	A	G	C	A	2




Figure 7. CpG island of HSP70 gene promoter and exon in Kampung Unggul Balitbangtan (KUB), Walik, and Kate Walik chickens

Table 5. Sequence base composition of HSP70 gene promoter and exon, and CpG island in Kampung Unggul Balitbangtan (KUB), Walik, and Kate Walik chickens

Breed	Sequence promoter and exon (bp)	Base composition (%)					
		A	T	G	C	A+T	G+C
Ref. chicken	787	23.8	18.2	32.7	25.3	42.0	58.0
KUB							
Haplotype 1	786	23.8	18.2	32.7	25.3	42.0	58.0
Haplotype 2	786	23.8	18.2	32.8	25.2	42.0	58.0
Haplotype 3	786	23.7	18.2	32.8	25.3	41.9	58.1
Walik							
Haplotype 1	786	23.8	18.2	32.7	25.3	42.0	58.0
Haplotype 2	786	23.8	18.2	32.8	25.2	42.0	58.0
Haplotype 3	786	23.7	18.2	32.8	25.3	41.9	58.1
Haplotype 4	786	23.7	18.2	32.9	25.2	41.9	58.1
Kate Walik							
Haplotype 2	786	23.8	18.2	32.8	25.2	42.0	58.0
CpG island 1	376	20.5	17.8	38.6	23.1	38.3	61.7

have potential to be used for marker-assisted selection (MAS) of reproductive traits in Pasundan cattle (Said & Putra, 2018).

In most organisms, the HSP70 gene does not usually have introns, is quite conserved, and, in many avian species, the protein coding regions have the same length i.e., 1,905 bp (Morimoto *et al.*, 1986; Mazzi *et al.*, 2003; Xia *et al.*, 2013; Zhang *et al.*, 2015). The complete coding-region sequences of HSP70 gene are available at NCBI for a number of avian species, including chicken (*G. gallus*, with accession numbers of J02579, AY143691, AY143692, and AY143693), guinea fowl (*Numida meleagris*, with accession number of AB096696), Japanese quail (*Coturnix japonica*, with accession number of

AB259847), duck (*Anas platyrhynchos*, with accession number of EU678246), and goose (*Anser cygnoides*, with accession number EU680475). The coding region of avian HSP70 gene generally starts with the standard start codon of ATG, ends with the stop codon of TAA, and codes for 633 amino acids.

Based on our analysis of the protein coding region (465 bp) of HSP70 genes in KUB, Walik, and Kate Walik chickens, four haplotypes were detected (Table 4). Mazzi *et al.* (2003) show that three hsp70 alleles (hsp70-1, hsp70-2, and hsp70-3) are found based on the polymorphisms at positions g.370A>G (identical to g.258A>G) and g.388C>G (identical to g.276C>G) in the protein coding region of the HSP70 genes of broiler and naked neck

chickens. We recognize that our H1 is 100% identical to the hsp70-3 allele in Pescoco Pelado 1 (with accession number of AY143692), H2 is 100% similar to the hsp70-2 allele in Hubbard Pettersen (with accession number of AY143693), and H3 is 100% identical to the hsp70-1 allele in Pescoco Pelado 1 (with accession number of AY143691). We find a new haplotype, H4, that is only found in Walik chickens in this study.

The mutation that is occurred at position g.370A>G produces the amino acid serine, while the mutation at position g.388C>G produces the amino acid proline. The two mutations have no changes to the original amino acid sequence of the HSP70 and are so-called silent mutations (Mazzi *et al.*, 2003). The silent mutations occurred in the coding region of the chicken HSP70 gene have been used as molecular markers for determining heat resistance in chickens (Mazzi *et al.*, 2003). Polymorphisms in the chicken HSP70 gene are detected by the PCR single-strand conformation polymorphism (PCR-SSCP) method (Tamzil *et al.*, 2013a; Liang *et al.*, 2016) and PCR restriction fragment length polymorphism (PCR-RFLP) analysis (Duangjinda *et al.*, 2017). Research on the HSP70 gene using the PCR-SSCP method has succeeded in identifying seven genotypes in Kampung Lombok chickens (AA, AB, AC, CC, AD, DD, and BC), six genotypes in Arabian chickens (AA, AB, AC, CC, AD, and BC), but only one genotype (DD) in commercial laying chickens (Tamzil *et al.*, 2013a). Based on the associations observed between chicken breeds and HSP70 gene genotypes and heat stress, native chickens appear to have the best heat resistance, while the most highly heat-tolerant genotype is AD (Tamzil *et al.*, 2013b). Liang *et al.* (2016) have shown the association between HSP70 gene polymorphisms detected by PCR-SSCP method with the acute temperature tolerance, growth, and egg production in Taiwanese local chickens. Only AA genotype (g.258A>G) shows a high tolerance level to acute temperatures (at 40 °C for 1 hour), without affecting the growth performance and egg production (Liang *et al.*, 2016). Thailand local and broiler chickens produce C and M loci, based on PCR-RFLP analysis with the restriction enzymes of *CfrI* and *MmeI* (New England Biolabs, Ipswich, MA) in coding region of the HSP70 gene, but only C1C1 and C1C2 genotypes that can be used as heat stress markers (Duangjinda *et al.*, 2017).

CpG is shorthand for 5'-C-phosphate-G-3'. CpG islands are regions with high frequency of CpG sites. The usual formal definition of CpG islands is a region with at least 200 bp, a GC percentage of greater than 50%, and observed-to-expected CpG ratio greater than 60% (Gardiner-Garden & Frommer, 1987). Many genes have CpG islands associated with the start of the gene (promoter regions). Methylation of CpG sites in the promoter of a gene may inhibit gene expression. In vertebrate, CpG islands are typically occurred at or near the transcription start site of genes. A cytosine base (C) followed immediately by a guanine base (G) is rare in vertebrate DNA, because the cytosines in such arrangement tend to be methylated. CpG islands are important because they represent areas of the genomes that have for some reasons been protected from mutating properties by methylation occurred through evolutionary time.

In our study, two of 33 CpG sites are found in the consensus promoter element of chicken HSP70 gene: SP 1 and HSEs (Figure 7). Gan *et al* (2013) detect methylation levels of core promoter transcription sites (SP 1 and HSEs). The results indicate that there is no correlation between the DNA methylation of the transcription factor CpG and HSP70 expression.

CONCLUSION

Haplotype 4 is a novel haplotype that is only found in Walik chicken. The appearance of H4 is caused by the mutation received in two specific sites (g.370A>G and g.388C>G). Walik chicken has a complete haplotype of HSP70 gene so that it can be used as a standard for heat stress genotyping in Indonesia local chickens.

CONFLICT OF INTEREST

We declare that there is no conflict of interest with any financial, personal, or other relationships with other people or organization related to the material discussed in the manuscript.

ACKNOWLEDGEMENT

The authors would like to thank and appreciate the Directorate General of Higher Education (DIKTI), The Ministry of Research, Technology, and Higher Education of The Republic of Indonesia for the financial support through BPPDN scholarship. We are grateful to Dr. Tike Sartika for permission to use KUB chickens.

REFERENCES

- Araujo, P. R., K. Yoon, D. Ko, A. D. Smith, M. Qiao, U. Suresh, S. C. Burns, & L. O. F. Penalva. 2012. Before it gets started: regulating translation at the 5'UTR [review]. *Comp. Funct. Genomics*. 2012:475731. <https://doi.org/10.1155/2012/475731>
- Archana, P. R., I. Aleena, P. Pragua, M. K. Vidya, A. P. A. Niyas, M. Bagath, G. Krishnan, A. Manimaran, V. Beena, E. K. Kurien, V. Sejian, & R. Bhatta. 2017. Role of heat shock proteins in livestock adaptation to heat stress. *J. Dairy Vet. Anim. Res.* 5:00127. <https://doi.org/10.15406/jdvar2017.05.00127>
- Akerfelt, M., R. I. Morimoto, & L. Sistonen. 2010. Heat shock factors: integrators of cell stress, development and lifespan. *Nat. Rev. Mol. Cell Biol.* 11:545-555. <https://doi.org/10.1038/nrm2938>
- Carabaño, M.J., M. Ramón, A. Menéndez-Buxadera, A. Molina & C. Diaz. 2019. Selecting for heat tolerance. *Animal Front.* 9:62-68. <https://doi.org/10.1093/af/vfy033>
- Chen, Z. Y., W. W. Zhang, J. K. Gan, L. N. Kong, X. Q. Zhang, D. X. Zhang, & Q. B. Luo. 2016. Genetic effect of an A/G polymorphism in the HSP70 gene on thermotolerance in chicken. *Genet. Mol. Res.* 15: gmr.15028271. <https://doi.org/10.4238/gmr.15028271>
- Duangjinda, M., S. Tunim, C. Duangdaen, & W. Boonkum. 2017. HSP70 genotypes and heat tolerance of commercial and native chickens reared in hot and humid conditions. *Braz. J. Poultry Sci.* 19:007-018. <https://doi.org/10.1590/1806-9061-2016-0245>
- Fathi, M.M., A. Galal, S. El-Safty, & M. Mahrous. 2013. Naked neck and frizzle genes for improving chickens raised

- under high ambient temperature: I. Growth performance and egg production. *World's Poult. Sci. J.* 69:813-832. <https://doi.org/10.1017/S0043933913000834>
- Fathi, M.M., A. Galal, S. El-Safy, & M. Mahrous.** 2014. Naked neck and frizzle genes for improving chickens raised under high ambient temperature: II. Blood parameters and immunity. *World's Poult. Sci. J.* 70:165-172. <https://doi.org/10.1017/S0043933914000142>
- Fujimoto, M., & A. Nakai.** 2010. The heat shock factor family and adaptation to proteotoxic stress. *FEBS J.* 277:4112-4125. <https://doi.org/10.1111/j.1742-4658.2010.07827.x>
- Gan, J.K., D.X. Zhang, D.I. He, X.Q. Zhang, Z.Y. Chen, & Q.B. Luo.** 2013. Promoter methylation negatively correlated with mRNA expression but not tissue differential expression after heat stress. *Genet. Mol. Res.* 12:809-819. <https://dx.doi.org/10.4238/2013.MARCH.15.1>
- Gan, J. K., L. Y. Jiang, L. N. Kong, X. Q. Zhang, & Q. B. Luo.** 2015. Analysis of genetic diversity of the heat shock protein 70 gene on the basis of abundant sequence polymorphisms in chicken breeds. *Genet. Mol. Res.* 14:1538-1545. <https://dx.doi.org/10.4238/2015.March.6.1>
- Gardiner-Garden, M., & M. Frommer.** 1987. CpG islands in vertebrate genomes. *J. Mol. Biol.* 196:261-282. [https://doi.org/10.1016/0022-2836\(87\)90689-9](https://doi.org/10.1016/0022-2836(87)90689-9)
- Hall, T.** 2011. Bioedit: An important software for molecular biology. *GERF Bull Biosciences.* 2:60-61.
- Leigh, J. W., & D. Bryant.** 2015. PopART: Full-feature software for haplotype network construction. *Methods Ecol. Evol.* 6:1110-1116. <https://doi.org/10.1111/2041-210X.12410>
- Leppek, K., R. Das, & M. Barna.** 2018. Functional 5'UTR mRNA structures in eukaryotic translation regulation and how to find them. *Nat. Rev. Mol. Cell. Biol.* 19:158-174. <https://doi.org/10.1038/nrm.2017.103>
- Liang, H. M., D. Y. Lin, Y. D. Hsuuw, T. P. Huang, H. L. Chang, C. Y. Lin, H. H. Wu, & K. H. Hung.** 2016. Association of heat shock protein 70 gene polymorphisms with acute thermal tolerance, growth, and egg production traits of native chickens in Taiwan. *Arch. Anim. Breed.* 59:173-181. <https://doi.org/10.5194/aab-59-173-2016>
- Mazzi, C. M., J. A. Ferro, M. I. T. Ferro, V. J. M. Savino, A. A. D. Coelho, & M. Macari.** 2003. Polymorphism analysis of the hsp70 stress gene in broiler chickens (*Gallus gallus*) of different breeds. *Gen. Mol. Biol.* 26:275-281. <https://doi.org/10.1590/S1415-47572003000300010>
- Mohamed, A. S. A., A. R. Lozovskiy, & A. M. A. Ali.** 2019. Strategies to combat the deleterious impacts of heat stress through feed restrictions and dietary supplementation (vitamins, minerals) in broilers. *J. Indonesian Trop. Anim. Agric.* 44:155-166. <https://doi.org/10.14710/jitaa.44.2.155-166>
- Morimoto, R. I., C. Hunt, S. Y. Huang, K. L. Berg, & S. S. Benerji.** 1986. Organization, nucleotide sequence, and transcription of the chicken HSP70 gene. *J. Biol. Chem.* 261:12692-12699.
- Nataamijaya, A. G.** 2010. Pengembangan potensi ayam lokal untuk menunjang peningkatan kesejahteraan petani. *Jurnal Litbang Pertanian* 29:131-138.
- Ng, C. S., P. Wu, J. Foley, A. Foley, M-L. McDonald, W-T Juan, C-J Huang, Y-T. Lai, W-S. Lo, C-F. Chen, S. M. Leal, H. Zhang, R. B. Widelitz, P. I Patel, W-H Li, & C-M. Chuong.** 2012. The chicken frizzle feather is due to an α -Keratin (KRT75) mutation that causes a defective rachis. *PLoS Genet.* 8:e1002748. <https://doi.org/10.1371/journal.pgen.1002748>
- Pelham, H. R. B.** 1982. A regulatory upstream promoter element in the *Drosophila* HSP70 heat-shock gene. *Cell.* 30:517-528. [https://doi.org/10.1016/0092-8674\(82\)90249-5](https://doi.org/10.1016/0092-8674(82)90249-5)
- Rayani, T. F., R. Mutia, & Sumiati.** 2017. Supplementation of zinc and vitamin E on apparent digestibility of nutrient, carcass traits, and mineral availability in broiler chickens. *Med. Pet.* 40:20-27. <https://doi.org/10.5398/medpet.2017.40.1.20>
- Said, S., & W. P. B. Putra.** 2018. Novel single nucleotide polymorphisms (SNPs) in the 5'UTR of bovine heat shock protein 70 (bHSP₇₀) gene and its association with service per conception (S/C) of pasundan cattle [short communication]. *Biodiversitas* 19:1622-1625. <https://doi.org/10.13057/biodiv/d190504>
- Sartika, T., S. Iskandar, & B. Tiesnamurti.** 2016. Sumberdaya Genetik Ayam Lokal Indonesia dan Prospek Pengembangannya. IAARD Press, Jakarta.
- Tamura, K., G. Stecher, D. Peterson, A. Filipski, & S. Kumar.** 2013. MEGA6: Molecular evolutionary genetics analysis version 6. *Mol. Biol. Evol.* 30:2725-2729. <https://doi.org/10.1093/molbev/mst197>
- Tamzil, M. H., R. R. Noor, P. S. Hardjosworo, W. Manalu, & C. Sumantri.** 2013a. Keragaman gen heat shock protein 70 pada ayam kampung, ayam arab dan ayam ras. *J. Vet.* 14:317-326.
- Tamzil, M. H., R. R. Noor, P. S. Hardjosworo, W. Manalu, & C. Sumantri.** 2013b. Acute heat stress responses of three lines of chickens with different heat shock protein (HSP)-70 genotypes. *Int. J. Poult. Sci.* 12:264-272.
- Tamzil, M. H.** 2014. Stres panas pada unggas: metabolisme, akibat dan upaya penanggulangannya. *Wartazoa* 24:57-66.
- Tkáčová, J., & M. Angelovičová.** 2012. Heat shock protein (HSPs): a review. *J. Anim. Sci. Biotechnol.* 45:349-353.
- Xia, M., J. Gan, Q. Luo, X. Zhang, & G. Yang.** 2013. Identification of duck HSP70 gene, polymorphism analysis and tissue expression under control and heat stress conditions. *Br. Poult. Sci.* 54:562-566. <https://dx.doi.org/10.1080/00071668.2013.819487>
- Zhang, W. W., X. Xiao, J. K. Gan, X. Q. Zhang, L. N. Kong, & Q. B. Luo.** 2015. Characterization of HSP70 and its expression in tissue: correlation with physiological and immune indices in goose (*Anser cygnoides*) serum. *Genet. Mol. Res.* 14:12288-12298. <https://doi.org/10.4238/2015.October.9.17>