

## Detection of Class 1 Integron Encoding Gene in Multidrug Resistance (MDR) *Citrobacter freundii* Isolated from Healthy Broiler Chicken

A. M. Witaningrum<sup>a</sup>, F. J. Wibisono<sup>b</sup>, D. A. Permatasari<sup>a</sup>, & M. H. Effendi<sup>a,\*</sup>

<sup>a</sup>Department of Veterinary Public Health, Faculty of Veterinary Medicine, Universitas Airlangga

<sup>b</sup>Department of Veterinary Public Health, Faculty of Veterinary Medicine, Universitas Wijaya Kusuma, Surabaya  
Jalan Kampus C Mulyorejo, FKH UNAIR, Surabaya 60115, Indonesia

\*Corresponding author: [mheffendi@yahoo.com](mailto:mheffendi@yahoo.com)

(Received 01-09-2020; Revised 18-01-2021; Accepted 16-02-2021)

### ABSTRACT

This study was aimed to find out that broiler chicken farms have problems with antibiotic resistance *Citrobacter freundii* and determined the prevalence and class 1 integron encoding gene. Multidrug resistance *Citrobacter freundii* was collected from broiler chicken among one hundred and sixty cloacal swab samples from 32 farms in Blitar for 3 months. The method of bacterial inoculation used MacConkey agar and biochemical test was conducted by IMViC and TSIA test. *Citrobacter freundii* for antibiotic sensitivity pattern was tested by disk diffusion, and the multidrug resistance encoding gene was tested by PCR. This study exposed 160 samples, and 13.75% (22/160) samples were positive of *Citrobacter freundii*. The antibiotic sensitivity pattern showed high resistances against ampicillin and erythromycin (77.27%), tetracycline (59.09%), trimetropim-sulfamethoxazole (50%), and streptomycin (22.72%). Isolates that were detected as multidrug resistance were continued with PCR testing to prove the existence of a class 1 integron encoding gene. Multidrug resistance *Citrobacter freundii* isolated from broiler chicken farms in Blitar were 81.82% (18/22), and were indicated that five were positive Class 1 Integron encoding gene. The results of this study showed that the prevalence and distribution of multidrug resistance *Citrobacter freundii* were high, so it can cause the spread of antimicrobial resistance to public health. Class 1 integron encoding gene was found 22.72% from multidrug resistance *Citrobacter freundii* by PCR. It was concluded that broiler chicken farms need assessment management to reduce and avoid multidrug resistance bacteria in animals and human. Therefore, the use of appropriate antibiotics is a good step to reduce the incidence of MDR in poultry.

**Keywords:** broiler chicken; *Citrobacter freundii*; Class 1 integron encoding gene; multidrug resistance; public health

### INTRODUCTION

Antibiotic is commonly used to prevent bacteria infection in poultry (Mehdi *et al.*, 2018). Poultry management applied more than one antibiotic to control bacteria (Roth *et al.*, 2019). Multi antibiotics uses for treatment caused resistance to bacteria and became a global public health problem (Prestinaci *et al.*, 2015). Antibiotics have been used for the treatment and prevention of disease in animals. Food of animal origin contains large amounts of antibiotic residues due to high antibiotic treatment, and this residue can be transmitted from animal products to humans (Landers *et al.*, 2012). Antibiotic resistance was found and detected at the Class 1 integrons encoding gene in poultry (Asgharpour *et al.*, 2018).

Integrons are gene to express encoding for antibiotic resistance (Kheiri *et al.*, 2016). Integrons are discovered on the transposons and contain inserted gene cassettes. The characterization of integrons is to harbor gene cassettes coding for resistance to antimicrobial (Domingues *et al.*, 2012; Permatasari *et al.*, 2020). Class 1 integrons are frequently reported with high prevalence in Gram

negative bacteria, including *Citrobacter freundii*. The class 1 integrons are found to be pathogen and made a world-wide significant bacterial infection in livestock (Karimi *et al.*, 2020; Hidayatullah *et al.*, 2020).

Enterobacteriaceae is frequently found in poultry (Projahn *et al.*, 2018), especially *Citrobacter freundii*. *Citrobacter freundii* is Gram negative, coccobacilli, motile using flagella and facultative anaerobic bacteria and commonly found in food, soil, water, and the intestines of animals and humans (Liu *et al.*, 2018). *Citrobacter freundii* existed in gastrointestinal and distributed food borne diseases of animals (Bai *et al.*, 2012; Prota *et al.*, 2015). Antibiotics inhibited the growth of bacteria, and treatment of multiple drugs caused resistance in bacteria infection. Multidrug resistance has been the global public problem (Liu *et al.*, 2018; Ansharieta *et al.*, 2020). Therefore, a detection method is needed at the molecular level to prove the spread of multidrug resistance (MDR) in poultry farms. The use of the Class 1 integron gene coding for the detection of MDR in *C. freundii* on poultry is a new step in the right direction.

This study was designed to evaluate the class 1 integron in broiler poultry farms and assessment of the multidrug resistance encoding gene amongst *Citrobacter freundii*. This study was designed to determine the possibility of emerge multidrug resistance caused by class 1 integron from bacteria in poultry farms.

## MATERIALS AND METHODS

### Ethical Clearance

Cloacal swabs were used in this study. Hence ethical clearance was not necessary. Cloacal swabs were collected from broiler chicken farms in Blitar, East Java province, Indonesia.

### Sample Isolation and Identification

This cross-sectional study was conducted between June and August 2019. One hundred and sixty cloacal swab samples were collected from broiler chicken farms in Blitar, Indonesia. The Sampling of the farm was based on less maintenance, sanitation, cleanliness, and hygiene management (Effendi *et al.*, 2018). The cloacal swab samples were cultured and incubated in MacConkey agar plates (Oxoid, Cheshire, UK) at 37°C for 24 h (Al Humam, 2016). The positive of *Citrobacter freundii* were indicated by lactose fermenting in MacConkey agar and colonies of bacteria obtain acquire pure subcultures (Liu *et al.*, 2017). Colonies of *Citrobacter freundii* were confirmed by biochemistry test. Indol test, Simmon Citrate Agar and TSIA were biochemistry tests to identify genus and species of *Citrobacter freundii* (Janda *et al.*, 1994; Nayar *et al.*, 2014).

### Antibiotic Sensitivity Test

Kirby-Bauer disc diffusion assay on medium Mueller-Hilton agar was done for testing antibiotic sensitivity (Nassar *et al.*, 2019; Effendi *et al.*, 2019). Erythromycin 15 µg, streptomycin (10 µg), tetracycline (30 µg), trimethoprim-sulfamethoxazole (25 µg), and ampicillin (10 µg) were antibiotics used to represent the antibiotic resistance according to Clinical and Laboratory Standards Institute recommendation. Antimicrobial susceptibility testing was carried out by measurement of inhibitory zone diameter formed at 37°C for 24 hours. (Clinical and Laboratory Standards Institute, 2017).

### Detection of Class 1 Integron Encoding Gene

*Citrobacter freundii* was revealed multidrug resistance using PCR method and analysis subtypes Class I Integron encoding gene identification. DNA isolation was done with the QIAamp® DNA mini kit (QIAGEN, Germany). The primers encoding the gene Class 1 Integron were hep 58 (TCATGGCTTGTTATGACTGT) and hep 59 (GTAGGGCTTATTATGCACGC). We used GoTaq Green mastermix (Promega, USA) and PCR condition with denaturation temperatures for 2 minutes at 94°C; extended denaturation at 94°C for 30 seconds;

annealing at 55°C for 45 seconds; extension at 72°C for 45 seconds; final extension at 72°C for 7 minutes, this reaction is carried out for 30 cycles. PCR results were confirmed by electrophoresis using 2% agarose gel (Invitrogen, USA), the amplicon using primer hep 58 and hep 59 was 200 bp (Singh *et al.*, 2017).

## RESULTS

Cloacal swab samples were collected from 160 chickens in broiler poultry farms in Blitar and the appearance of multidrug resistance (MDR) *Citrobacter freundii* was detected. The sample was obtained 13.75% (22/160) *Citrobacter freundii* and 81.82% (18/22) multidrug resistance cases in broiler poultry farms.

The antibiotic sensitivity test showed that the highest multidrug resistance *Citrobacter freundii* to ampicillin and erythromycin (77.27%), tetracycline (59.09%), trimetropim-sulfamethoxazole (50.00%), and streptomycin (22.72%) (Table 1). The most multidrug resistance showing sensitive to antibiotic is streptomycin (77.27%). Antibiotics are commonly used to affect sensitive bacteria, but if used inappropriately, they can lead to antibiotic resistance (Singh *et al.*, 2017).

The amplification Class 1 Integron of *Citrobacter freundii* from 22 cloacal swab samples of broiler poultry was positive for 200 bp using the primer hep 58 and hep 59 (Nagachinta *et al.*, 2009). The results indicated that five were positive Class 1 Integron encoding genes (Table 2).

The results showed that several resistance antibiotics of *Citrobacter freundii* are related to the appearance of the Class 1 Integron. Figure 1 shows *Citrobacter freundii* are resistance to several antibiotics. Figure 2 shows that multidrug resistance to antibiotics. The multidrug resistance *Citrobacter freundii* by antibiotic susceptibility testing was confirmed 18/22 (81.82%), and Class 1 Integron encoding gene positive by PCR testing was showed 5/22 (22.72%) (Figure 3).

## DISCUSSION

The study discovered contamination of *Citrobacter freundii* caused pathogen and foodborne diseases in broiler farms (Aminharati *et al.*, 2019). The multidrug resistance to several antibiotics was reported from isolated 160 cloacal swab samples. The major public health problem has many cases especially higher prevalence of

Table 1. Antibiotic susceptibility profile of *Citrobacter freundii* in broiler chicken farms in Blitar, Indonesia

Antibiotics	Resistance (%)	Intermediate (%)	Sensitive (%)
Ampicillin (AMP)	17 (77.27%)	1 (4.55%)	4 (18.18%)
Erythromycin (E)	17 (77.27%)	3 (13.64%)	2 (9.09%)
Tetracycline (TE)	13 (59.09%)	0	9 (40.91%)
Trimethoprim-sulfamethoxazole (SXT)	11 (50.00%)	0	11 (50.00%)
Streptomycin (S)	5 (22.73%)	0	17 (77.27%)

Table 2. Samples result of multidrug resistant (MDR) and Class 1 Integron cases on *Citrobacter freundii*

Locations	Sample of <i>Citrobacter freundii</i>	Positive <i>Citrobacter freundii</i>	Positive MDR	Positive Class 1 Integron by PCR
Farm 3	B3an, <b>B3bn</b> , B3cn	3 (13.64%)	2 (9.09%)	1 (4.55%)
Farm 4	B4bn, B4dn	2 (9.09%)	0	0
Farm 7	<b>B7cn</b>	1 (4.55%)	1 (4.55%)	0
Farm 8	<b>B8an</b> , <b>B8en</b>	2 (9.09%)	2 (9.09%)	0
Farm 11	B11en	1 (4.55%)	1 (4.55%)	1 (4.55%)
Farm 14	B14an, B14bn	2 (9.09%)	2 (9.09%)	2 (9.09%)
Farm 26	<b>B26bn</b>	1 (4.55%)	1 (4.55%)	0
Farm 27	<b>B27an</b> , <b>B27en</b>	2 (9.09%)	2 (9.09%)	0
Farm 28	<b>B28cn</b> , B28dn	2 (9.09%)	1 (4.55%)	0
Farm 30	<b>B30an</b> , <b>B30dn</b> , <b>B30en</b>	3 (13.64%)	3 (13.64%)	0
Farm 31	<b>B31an</b> , B31cn, <b>B31en</b>	3 (13.64%)	3 (13.64%)	1 (4.55%)

Note: Bold code= positive result of MDR *Citrobacter freundii*; Italic code= positive result of Class 1 Integron gene.




Figure 1. Antibiotic susceptibility profile of *Citrobacter freundii* from broiler chicken farms in Blitar, Indonesia with result was multidrug resistant (MDR). The antibiotic were Ampicillin (AMP), Erythromycin (E), Tetracycline (TE), Trimethroprim-sulfamethoxazole (SXT), and Streptomycin (S). The samples were incubated at 37°C for 24 hours.

Note: AMP, E, SXT, and S were resistant to *Citrobacter freundii*; TE was sensitive. Therefore, this isolate was MDR due to more than three antibiotics resistant.

multidrug resistant bacterial pathogens. Integrons are related to multidrug resistance in many cases of bacterial diseases (Akrami *et al.*, 2019; Krauland *et al.*, 2009). Integrons were recognized and detected first time in Gram-negative bacteria (Deng *et al.*, 2015; Domingues *et al.*, 2012; Pormohammad *et al.*, 2019).

Organisms that are resistant to three or more classes of antimicrobials are referred to as multidrug resistant (MDR) (Magiorakos *et al.*, 2012; Wibisono *et al.*, 2020). One method frequently used by various researchers to characterize organisms as MDR is based on the results of *in vitro* antimicrobial susceptibility testing (Kallen *et al.*, 2010). The most commonly used definition is for Gram-negative bacteria that are resistant to three or more classes of antimicrobials (Gould, 2008, Kristianingtyas *et al.*, 2020). The variability of this definition is provided in the comprehensive MDR review




Figure 2. Percentage of antibiotic resistance on *Citrobacter freundii*. AMP (Ampicillin), E (Erythromycin), TE (Tetracycline), SXT (Trimethroprim-sulfamethoxazole), and S (Streptomycin).

(Falagas *et al.*, 2006), which is used by some researchers as a reference that a large number of studies do not propose a specific definition for MDR.

There were 18 MDR of *Citrobacter freundii* isolates in this study, shown in Table 2, although 5 isolates contain Class 1 integrons gene of *Citrobacter freundii* in this study. Other isolates are shown negative of Class 1 integrons gene because globally pathogens of Class 1 integrons discovered on the chromosomes of environmental bacteria. Four classes of integron have been described, each of which codes for a distinct but related integrase enzyme. Class 1 integrons are the most widely studied (Rosser & Young, 1999). Class 1 integrons are bounding in conjugative plasmids, transposons conjugative plasmids, and spreading by lateral gene transfer. Therefore, Class 1 integrons have overspread to almost all species of Gram-negative pathogens. Class 1 integron is a major competence in the global spread of multidrug resistance and important to recover management of the farm to decrease the prevalence of antibiotic resistance and improving the balance of healthy life.

Consumption of antibiotics increased antibiotic resistance containing important MDR organisms in poultry. These MDR organisms can be transmitted to humans




Figure 3. Class 1 Integron gene of *Citrobacter freundii* from broiler chicken farms in Blitar, Indonesia. MRK (marker), K+ (positive control) was used *Escherichia coli* from patients of multidrug resistance (MDR), K- (negative control) was used *Pseudomonas aeruginosa* ATCC 27853. A) B3cn, B11en, B14an, B14bn were positive result of Class 1 Integron; B) B31cn was positive result of Class 1 Integron.

through direct contact or consumption, and *C. freundii* causes economic losses and levels of food contamination. Therefore antibiotic-resistant genes have the potential to spread to other populations. The abundant use of antibiotics in poultry farming has been linked to treatment failure and the development of antibiotic resistance itself.

Integrations are genes encoding multidrug resistance and reduction of treatment for bacterial infections (Mostafa *et al.*, 2015). The classification of integrons is based on differences in the gene structure of integrases (Cury *et al.*, 2015). The spread of Class 1 integrons have been found commonly in Gram-negative bacteria and showed the existence of Class 1 integrons of *Citrobacter freundii* in this study. The appearance result of amplification class 1 integron from positive isolates were 22.72%. The amplification using primer hep 58 and hep 59 for 200bp represent class 1 integron of *Citrobacter freundii*. Multidrug resistance was reported to be spread by Class 1 integron genes in Gram-negative bacteria.

The multidrug resistance nature of this isolate can be explained by the fact that it is mediated by plasmids carrying multiresistant genes and by transposons, and by integrons that are easily transferred to other bacteria, not necessarily of the same species (Widodo *et al.*, 2020). Bacteria with various resistance to antibiotics are widespread in animals and the environment (Kwoji *et al.*, 2019, Riwu *et al.*, 2020). Recent surveys from China (Gao *et al.*, 2015), Thailand (Runcharoen *et al.*, 2017), and Indonesia (Effendi *et al.*, 2018), have illustrated an alarming trend regarding resistance among broad-spectrum beta lactamase-producing organisms (ESBLs), which are also multidrug-resistant isolated from animals and the environment (Wibisono *et al.*, 2020).

Farm animals are frequently found to be infected by pathogenic bacteria and the infection can be transferred from animals to humans, especially *Citrobacter freundii* (Zhou *et al.*, 2019; Liu *et al.*, 2017). The treatment

to pathogenic bacteria used large amounts of antibiotics to reduce attack infection and caused resistance to multiple drugs. Multidrug resistance bacteria are most important to observation concern because influence public health problem.

## CONCLUSION

*Citrobacter freundii* was discovered in broiler chicken. This study showed that multidrug resistance *Citrobacter freundii* was confirmed 18/22 (81.82%), and Class 1 Integron encoding gene positive by PCR testing was 5/22 (22.72%). The result revealed multidrug resistance Gram negative bacteria emerge in poultry farms and need evaluation management plans to prevent the transfer of multidrug resistance bacteria from poultry to humans and the environment.

## CONFLICT OF INTEREST

The authors declare that there is no conflict of interest with any financial, personal, or other relationships with other people or organization related to the material discussed in the manuscript.

## ACKNOWLEDGEMENT

This research was in part funded by the Direktorat Riset dan Pengabdian Masyarakat, Deputy Bidang Penguatan Riset dan Pengembangan Kementerian Riset dan Teknologi/ Badan Riset dan Inovasi Nasional, Indonesia in fiscal year 2020 with grant number: 756/UN3.14/PT/2020.

## REFERENCES

Akrami, F., M. Rajabnia, & A. Pournajaf. 2019. Resistance integrons; A mini review. *Caspian J. Intern. Med.* 10:370-376.

- Aminharati, F., M. H. Ehrampoush, S. M. Dallal, S. M., M. Yaseri, D. A. Tafti, & Z. Rajabi. 2019. *Citrobacter freundii* Foodborne disease outbreak related to environmental conditions in Yazd Province, Iran. Iran J. Public Health. 48:1099-1105. <https://doi.org/10.18502/ijph.v48i6.2919>
- Ansharieta, R., M. H. Effendi, & H. Plumeriastuti. 2020. Detection of multidrug-resistant (MDR) *Escherichia coli* isolated from raw milk in East Java Province, Indonesia. Indian J. Forensic Med. Toxicol. 14:4303-4307
- Asgharpour, F., S. Mahmoud, A. Marashi, & Z. Moulana. 2018. Molecular detection of class 1, 2 and 3 integrons and some antimicrobial resistance genes in *Salmonella infantis* isolates. Iran J. Microbiol. 10:104-110.
- Bai, L., S. Xia, R. Lan, L. Liu, C. Ye, Y. Wang, D. Jin, Z. Cui, H. Jing, Y. Xiong, X. Bai, H. Sun, J. Zhang, L. Wang, & J. Xu. 2012. Isolation and characterization of cytotoxic, aggregative *Citrobacter freundii*. PLoS ONE 7:e33054. <https://doi.org/10.1371/journal.pone.0033054>
- Clinical and Laboratory Standards Institute. 2017. Performances Standards for Antimicrobial Susceptibility Testing. 27<sup>th</sup> ed. CLSI supplement M100. Clinical and Laboratory Standards Institute, Wayne.
- Cury, J., T. Jové, M. Touchon, B. Néron, & E. P. Rocha. 2016. Identification and analysis of integrons and cassette arrays in bacterial genomes. Nucleic Acids Res. 44:4539-4550. <https://doi.org/10.1093/nar/gkw319>
- Dehkordi, M. K., M. Halaji, & S. Nouri. 2020. Prevalence of class 1 integron in *Escherichia coli* isolated from animal sources in Iran: a systematic review and meta-analysis. Trop. Med. Health. 48:1-7. <https://doi.org/10.1186/s41182-020-00202-1>
- Deng, Y., X. Bao, L. Ji, L. Chen, J. Liu, J. Miao, D. Chen, H. Bian, Y. Li, & G. Yu. 2015. Resistance integrons: class 1, 2 and 3 integrons. Ann. Clin. Microbiol. Antimicrob. 14:1-11. <https://doi.org/10.1186/s12941-015-0100-6>
- Domingues, S., G. J. da Silva, & K. M. Nielsen. 2012. Integrons: Vehicles and pathways for horizontal dissemination in bacteria. Mobile genetic elements 2:211-223. <https://doi.org/10.4161/mge.22967>
- Effendi, M. H., I.G. Bintari, E. B. Aksono, & I. P. Hermawan. 2018. Detection of blaTEM gene of *Klebsiella pneumoniae* isolated from swab of food-producing animals in East Java. Trop. Anim. Sci. J. 41:174-178. <https://doi.org/10.5398/tasj.2018.41.3.174>
- Effendi M.H., N. Harijani, S. M. Yanestria & P. Hastutiek. 2018. Identification of shiga toxin-producing *Escherichia coli* in raw milk samples from dairy cows in Surabaya, Indonesia. Philipp. J. Vet. Med. 55:109-114.
- Effendi, M.H., N. Harijani, Budiarto, N. P. Triningtya, W. Tyasningsih, & H. Plumeriastuti. 2019. Prevalence of pathogenic *Escherichia coli* Isolated from Subclinical Mastitis in East Java Province, Indonesia. Indian Vet. J. 96:22-25. <https://doi.org/10.13057/biodiv/d220137>
- Falagas, M. E., P. K. Koletsis, & I. A. Bliziotis. 2006. The diversity of definitions of multidrug-resistant (MDR) and pandrug-resistant (PDR) *Acinetobacter baumannii* and *Pseudomonas aeruginosa*. J. Med. Microbiol. 55:1619-1629. <https://doi.org/10.1099/jmm.0.46747-0>
- Gould, I. M. 2008. The epidemiology of antibiotic resistance. Int. J. Antimicrob. Agents. 32 (suppl 1): 2-9. <https://doi.org/10.1016/j.ijantimicag.2008.06.016>
- Hidayatullah, A. R., M. H. Effendi, H. Plumeriastuti, F. M. Wibisono, E.B. Hartadi, & E. D. Sofiana. 2020. A Review of the opportunistic pathogen *Citrobacter freundii* in piglets post weaning : Public Health Importance. Sys. Rev. Pharm. 11:767-773.
- Humam, N. A. 2016. Special biochemical profiles of *Escherichia coli* strains isolated from humans and camels by the VITEK 2 automated system in Al-Ahsa, Saudi Arabia. Afr. J. Microbiol. Res.10: 783-790. <https://doi.org/10.5897/AJMR2016.8047>
- Janda, J. M., S. L. Abbott, W. K. W. Cheung, & D. F. Hanson. 1994. Biochemical identification of *Citrobacteria* in the clinical laboratory. J. Clin. Microbiol. 32: 1850-1854. <https://doi.org/10.1128/JCM.32.8.1850-1854.1994>
- Kallen, A. J., A. I. Hidron, J. Patel, & A. Srinivasan. 2010. Multidrug resistance among gram-negative pathogens that caused healthcare-associated infections reported to the National Healthcare Safety Network, 2006-2008. Infect. Control Hosp. Epidemiol. 31:528-531. <https://doi.org/10.1086/652152>
- Kheiri, R. & L. Akhtari. 2016. Antimicrobial resistance and integron gene cassette arrays in commensal *Escherichia coli* from human and animal sources in IRI. Gut pathog. 8:1-10. <https://doi.org/10.1186/s13099-016-0123-3>
- Krauland, M. G., J. W. Marsh, D. L. Paterson, & L. H. Harrison. 2009. Integron-mediated multidrug resistance in a global collection of nontyphoidal *Salmonella enterica* isolates. Emerg. Infect. Dis. 15:388-396. <https://doi.org/10.3201/eid1503.081131>
- Kristianingtyas, L., M. H. Effendi, W. Tyasningsih, & F. Kurniawan. 2020. Genetic identification of blaCTX-M gene and blaTEM gene on Extended Spectrum Beta Lactamase (ESBL) producing *Escherichia coli* from dogs. Indian Vet. J. 97:17-21
- Kwoji, I. D., J. A. Musa, N. Daniel, D. L. Mohzo, A. A. Bitrus, A. A. Ojo, & K. U. Ezema. 2019. Extended-spectrum beta-lactamase-producing *Escherichia coli* in chickens from small-scale (backyard) poultry farms in Maiduguri, Nigeria. Int. J. One Health. 5:26-30. <https://doi.org/10.14202/IJOH.2019.26-30>
- Landers, T. F., B. Cohen, T. E. Wittum, & E. L. Larson. 2012. A review of antibiotic use in food animals: perspective, policy, and potential. Public Health Rep. 127:4-22. <https://doi.org/10.1177/003335491212700103>
- Liu, L., D. Chen, L. Liu, R. Lan, S. Hao, W. Jin, H. Sun, Y. Wang, Y. Liang, & J. Xu. 2018. Genetic diversity, multidrug resistance, and virulence of *Citrobacter freundii* from diarrheal patients and healthy individuals. Front. Cell. Infect. Microbiol. 8:1-10 <https://doi.org/10.3389/fcimb.2018.00233>
- Liu, L. H., N. Y. Wang, A. Y. J. Wu, C. C. Lin, C. M. Lee, & C. P. Liu. 2018. *Citrobacter freundii* bacteremia: Risk factors of mortality and prevalence of resistance genes. J. Microbiol. Immunol. Infect. 51:565-572. <https://doi.org/10.1016/j.jmii.2016.08.016>
- Liu, L., R. Lan, L. Liu, Y. Wang, Y. Zhang, Y. Wang, & J. Xu. 2017. Antimicrobial resistance and cytotoxicity of *Citrobacter* spp. in Maanshan Anhui Province, China. Front. Microbiol. 8:1357. <https://doi.org/10.3389/fmicb.2017.01357>
- Magiorakos, A.P., A. Srinivasan, R. B. Carey, Y. Carmeli, M. E. Falagas, C. G. Giske, S. Harbarth, J. F. Hindler, G. Kahlmeter, B. O. Liljequist, D. L. Paterson, L. B. Rice, J. Stelling, M. J. Struelens, A. Vatopoulos, J. T. Weber, & D. L. Monnet. 2012. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. Clin. Microbiol. Infect. 18:268-281. <https://doi.org/10.1111/j.1469-0691.2011.03570.x>
- Mehdi, Y., M. P. L. Montminy, M. L. Gaucher, Y. Chorfi, G. Suresh, T. Rouissi, S. K. Brar, C. Côté, A. A. Ramirez, & S. Godbout. 2018. Use of antibiotics in broiler production: Global impacts and alternatives. Anim. Nutr. 4:170-178. <https://doi.org/10.1016/j.aninu.2018.03.002>
- Mostafa, M., S. D. Siadat, F. Shahcheraghi, F. Vaziri, A. J. Nejad, J. V. Yousefi, B. Rajaei, E. H. Mood, N. E. Zadeh, A. Moshiri, S. A. S. Siamdoust, & M. Rahbar. 2015. Variability in gene cassette patterns of class 1 and

- 2 integrons associated with multi drug resistance patterns in *Staphylococcus aureus* clinical isolates in Tehran-Iran. BMC microbial. 15:152. <https://doi.org/10.1186/s12866-015-0488-3>
- Nagachinta, S. & J. Chen.** 2009. Integron-mediated antibiotic resistance in Shiga Toxin-producing *Escherichia coli*. J. Food Prot. 72:21-27. <https://doi.org/10.4315/0362-028X-72.1.21>
- Nassar, M. S. M., W. A. Hazzah, & W. M. K. Bakr.** 2019. Evaluation of antibiotic susceptibility test results: how guilty a laboratory could be?. J. Egypt Public Health Assoc. 94:4. <https://doi.org/10.1186/s42506-018-0006-1>
- Nayar, R., I. Shukla, & A. Sultan.** 2014. Epidemiology, prevalence and identification of *Citrobacter* species in clinical specimens in a tertiary care hospital in India. International Journal of Scientific and Research Publications 4:4.
- Permatasari, D. A., A. M. Witaningrum, F. J. Wibisono, & M. H. Effendi.** 2020. Detection and prevalence of Multidrug Resistant *Klebsiella pneumoniae* strains isolated from poultry farms in Blitar, Indonesia. Biodiversitas 21:4642-4647. <https://doi.org/10.13057/biodiv/d211024>
- Pormohammad, A., R. Pouriran, H. Azimi, & M. Goudarzi.** 2019. Prevalence of integron classes in Gram-negative clinical isolated bacteria in Iran: A systematic review and meta-analysis. Iran J. Basic Med. Sci. 22:118-127.
- Prestinaci, F., P. Pezzotti, & A. Pantosti.** 2015. Antimicrobial resistance: A global multifaceted phenomenon. Pathog. Glob. Health. 109:309-318. <https://doi.org/10.1179/2047773215Y.00000000030>
- Projahn, M., E. Pacholewicz, E. Becker, G. C. Carreira, N. Bandick, & A. Kaesbohrer.** 2018. Reviewing interventions against *Enterobacteriaceae* in Broiler processing: Using old techniques for meeting the new challenges of ESBL *E. coli*?. Biomed Res. Int. 2018:7309346. <https://doi.org/10.1155/2018/7309346>
- Prota, M. A., A. P. Sandoval, M. G. Clemente, R. Fernández, & P. Casan.** 2015. Community-acquired pneumonia and empyema caused by *Citrobacter koseri* in an immunocompetent patient. Case Rep. Pulmonol. 2015:1-6. <https://doi.org/10.1155/2015/670373>
- Riwu, K. H. P., M. H. Effendi, & F. A. Rantam.** 2020. A review of extended Spectrum  $\beta$ -Lactamase (ESBL) producing *Klebsiella pneumoniae* and Multidrug Resistant (MDR) on companion animals. Sys. Rev. Pharm. 11:270-277.
- Rosser, S. J. & Young, H. K.** 1999. Identification and characterization of class 1 integrons in bacteria from an aquatic environment. J. Antimicrob. Chemother. 44:11-18. <https://doi.org/10.1093/jac/44.1.11>
- Roth, N., A. Käsbohrer, S. Mayrhofer, U. Zitz, C. Hofacre, & K. J. Domig.** 2019. The application of antibiotics in broiler production and the resulting antibiotic resistance in *Escherichia coli*: A global overview. Poult. Sci. 98:1791-1804. <https://doi.org/10.3382/ps/pey539>
- Runcharoen, C., K. E. Raven, S. Reuter, T. Kallonen, S. Paksanont, & J. Thammachote.** 2017. Whole-genome sequencing of ESBL producing *Escherichia coli* isolated from patients, farm waste and canals in Thailand. Gen. Med. 9:81-93. <https://doi.org/10.1186/s13073-017-0471-8>
- Singh, T., S. Das, V. G. Ramachandran, S. Wani, D. Shah, K. A. Maroof, & A. Sharma.** 2017. Distribution of integrons and phylogenetic groups among enteropathogenic *Escherichia coli* isolates from children <5 years of age in Delhi, India. Front. Microbiol. 8:561. <https://doi.org/10.3389/fmicb.2017.00561>
- Wibisono, F. J., B. Sumiarto, T. Untari, M. H. Effendi, D. A. Permatasari, & A. M. Witaningrum.** 2020. The presence of Extended Spectrum Beta-Lactamase (ESBL) producing *Escherichia coli* on layer chicken farms in Blitar area, Indonesia. Biodiversitas 21: 2667-2671. <https://doi.org/10.13057/biodiv/d210638>
- Wibisono, F.J., B. Sumiarto, T. Untari, M.H. Effendi, D.A. Permatasari, & A.M. Witaningrum.** 2020. Short Communication: Pattern of antibiotic resistance on extended-spectrum beta-lactamases genes producing *Escherichia coli* on laying hens in Blitar, Indonesia. Biodiversitas 21: 4631-4635. <https://doi.org/10.13057/biodiv/d211022>
- Widodo, A., M. H. Effendi, & A. R. Khairullah.** 2020. Extended-spectrum beta-lactamase (ESBL)-producing *Escherichia coli* from livestock. Sys. Rev. Phar. 11: 382-392.
- Zhou, W., Q. Chen, C. Qian, K. Shen, X. Zhu, D. Zhou, W. Lu, Z. Sun, H. Liu, K. Li, T. Xu, Q. Bao, & J. Lu.** 2019. *In Vitro* susceptibility and florfenicol resistance in *Citrobacter* isolates and whole-genome analysis of multidrug-resistant *Citrobacter freundii*. Int. J. Genomics. 2019:1-15. <https://doi.org/10.1155/2019/7191935>